New forests and Kyoto Protocol carbon accounting: A case study in central Italy

Alessandro Alivernini^{a,*}, Anna Barbati^a, Paolo Merlini^b, Francesco Carbone^a,
Piermaria Corona^b

^a University of Tuscia—Department for Innovation in Biological, Agro-food and Forest Systems (DIBAF), Via S. Camillo De Lellis snc, 01100 Viterbo, Italy

^b Consiglio per la ricerca e l'analisi dell'economia agraria—Forestry Research Center (CREA-SEL), Viale Santa Margherita 80, 52100 Arezzo, Italy

ARTICLE INFO

Article history:

Received 16 July 2015

Received in revised form 1 October 2015

Accepted 3 November 2015

Available online xxx

Keywords:

Afforestation

Carbon sequestration

Kyoto Protocol

Airborne laser scanning

CO2Fix

ABSTRACT

Forest expansion into formerly agricultural land plays a relevant role for generating carbon sinks eligible under afforestation activities in the Kyoto Protocol (KP) carbon accounting framework. This paper provides a methodological framework for the multi-temporal assessment of carbon accumulated in these new forest stands, at the reference years of the KP commitment periods: remote sensing analyses, integrating optical imagery and airborne laser scanning data with field survey, are combined with the carbon simulation model CO2FIX to project carbon changes. The approach is applied to a case study in the central Apennines (Italy) to assess the carbon sink associated to these new forests at the years 2008, 2012 and 2020. The approach is based on forest cover change detection and growing stock mapping, while the multi-temporal carbon sink analysis is performed by means of the CO2FIX model. In the considered test area, the results highlight a compound annual rate of forest expansion equal to 0.26% for the period 1990–2012. The overall carbon stored by the new stands is 21.1 Mg ha^{-1} (relative standard error less than 10%) at the year 2012, projected to reach 33.5 Mg ha^{-1} at the year 2020.

© 2015 Elsevier B.V. All rights reserved.

1. Introduction

Countries ratifying the United Nations Framework Convention on Climate Change (UNFCCC) Kyoto Protocol (KP) are given the option to meet part of their reduction requirements through the carbon credits generated by afforestation activities after 1990 (Article 3.3). The protocol allows sinks associated with woodland creation in abandoned farmland to be included in the national carbon accounting system to offset emissions, provided that forest expansion can be considered a direct human-induced conversion of land that has not been forested for a period of at least 50 years (UNFCCC, 2001). The possibility to include afforestation as permissible offset activity holds also for the second commitment period of the KP (UNFCCC, 2011).

Italy has expressed a formal position to include the carbon sinks generated by forest expansion as afforestation activities; such a widespread woodland creation phenomenon is regarded human-induced, as it is a natural dynamic that would not otherwise occur

without intentional agricultural land abandonment. This phenomenon, widely observed in rural landscapes in Italy (Barbati et al., 2013; Cimini et al., 2013; Corona et al., 2008; Piussi and Pettenella, 2000), is common to many countries of the global north (Beilin et al., 2014). Yet, only few studies have quantitatively analyzed the implications of forest expansion in terms of creation of new carbon sinks (Acosta et al., 2005; Conti and Fagarazzi, 2004; Gils et al., 2008; Mazzoleni et al., 2004).

To bridge this knowledge gap, this paper addresses as a main scientific question the experimentation of a methodological approach, founded on the integration of forest mapping and inventory (Corona, 2010; McRoberts, 2010) with carbon modelling, to assess carbon accumulated in forest stands naturally originated by forest expansion, at the reference years of KP commitment periods (2008, 2012, 2020). Distinctively, simple remote sensing analyses are combined with the carbon simulation model CO2FIX (Masera et al., 2003) to project carbon changes (Fig. 1). The approach allows wall-to-wall mapping of above ground carbon stock, suitable to support decision-making at various administrative levels, from municipality up to subnational scale.

Under such a perspective, the proposed approach has been tested in a case study in central Italy. The experimental framework builds on three steps: (1) forest cover change detection to quantify natural afforestation in the period 1990–2012 by means of optical

* Corresponding author.

E-mail addresses: alivernini@unitus.it (A. Alivernini), barbati.sisfor@unitus.it (A. Barbati), paolo.merlini@entecra.it (P. Merlini), fcabone@unitus.it (F. Carbone), piermaria.corona@entecra.it (P. Corona).