

Knowledge gaps about mixed forests: What do European forest managers want to know and what answers can science provide?

Lluís Coll^{a,b,c,*}, Aitor Ameztegui^{b,c}, Catherine Collet^d, Magnus Löf^e, Bill Mason^f, Maciej Pach^g, Kris Verheyen^h, Ioan Abrudaniⁱ, Anna Barbat^j, Susana Barreiro^k, Kamil Bielak^l, Andrés Bravo-Oviedo^{m,n}, Barbara Ferrari^j, Zoran Govedar^o, Jiri Kulhavy^p, Dagnija Lazdina^q, Marek Metslaid^r, Frits Mohren^s, Mário Pereira^t, Sanja Peric^u, Ervin Rasztovits^v, Ian Short^w, Peter Spathelf^x, Hubert Sterba^y, Dejan Stojanovic^z, Lauri Valsta^{aa}, Tzvetan Zlatanov^{ab}, Quentin Ponette^{ac}

^a Department of Agriculture and Forest Engineering (EAGROF), University of Lleida, 25198 Lleida, Spain

^b Forest Sciences Centre of Catalonia (CTFC), Solsona 25280, Spain

^c CREAM, Centre for Ecological Research and Forestry Applications, Autonomous University of Barcelona, 08193 Cerdanyola del Vallès, Spain

^d LERFoB, UMR 1092, INRA-AgroParisTech, 54280 Champenoux, France

^e Southern Swedish Forest Research Centre, Swedish University of Agricultural Science, 230 53 Alnarp, Sweden

^f Forest Research, Northern Research Station, EH25 9SY Midlothian, Scotland, UK

^g Department of Silviculture, University of Agriculture in Krakow, 31-425 Krakow, Poland

^h Forest and Nature Lab, Ghent University, 9090 Melle-Gontrode, Belgium

ⁱ Faculty of Silviculture and Forest Engineering, Transilvania University of Brasov, 500123 Brasov, Romania

^j Department for the Innovation in Biological, Agrofood and Forest Systems, University of Tuscia, 01100 Viterbo, Italy

^k Forest Research Centre (CEF), School of Agriculture, University of Lisbon, 1349-017 Lisbon, Portugal

^l Department of Silviculture, Warsaw University of Life Sciences, 02-878 Warsaw, Poland

^m INIA-CIFOR, 28040 Madrid, Spain

ⁿ Sustainable Forest Management Research, Institute University of Valladolid & INIA, 34004 Palencia, Spain

^o Faculty of Forestry, University in Banja Luka, Banja Luka 78000, Bosnia and Herzegovina

^p Mendel University of Brno, Faculty of Forestry and Wood Technology, 61300 Brno, Czech Republic

^q Latvian State Forest Research Institute Silava, 2169 Salaspils, Latvia

^r Institute of Forestry and Rural Engineering, Estonian University of Life Sciences, 51014 Tartu, Estonia

^s Forest Ecology and Forest Management, Wageningen University of Environmental Sciences, 6708 Wageningen, The Netherlands

^t Centre for Research and Technology of Agro-Environment and Biological Sciences (CITAB), University of Trás-os-Montes and Alto Douro (UTAD), 5000-801 Vila Real, Portugal

^u Croatian Forest Research Institute, 10450 Jastrebarsko, Croatia

^v Institute of Environmental and Earth Sciences, University of West Hungary, 9400 Sopron, Hungary

^w Teagasc Forestry Development Department, Teagasc Ashtown Food Research Centre, Ashtown, Dublin 15, Ireland

^x Faculty of Forest and Environment, University for Sustainable Development Eberswalde, 16225 Eberswalde, Germany

^y Department of Forest and Soil Sciences, BOKU University of Natural Resources and Life Sciences, 1190 Vienna, Austria

^z Institute of Lowland Forestry and Environment, University of Novi Sad, 21000 Novi Sad, Serbia

^{aa} University of Helsinki, Department of Forest Sciences, 00014 Helsinki, Finland

^{ab} Forest Research Institute, 1756 Sofia, Bulgaria

^{ac} UCL, Earth and Life Institute, Environmental Sciences, 1348 Louvain-la-Neuve, Belgium

* Corresponding author at: Department of Agriculture and Forest Engineering (EAGROF), University of Lleida, Av. Alcalde Rovira Roure 191, 25198 Lleida, Spain.
E-mail address: lluis.coll@eagrof.udl.cat (L. Coll).

Abstract

Research into mixed-forests has increased substantially in the last decades but the extent to which the new knowledge generated meets practitioners' concerns and is adequately transmitted to them is unknown. Here we provide the current state of knowledge and future research directions with regards to 10 questions about mixed-forest functioning and selected by a range of European forest managers during an extensive participatory process. The set of 10 questions were the highest ranked questions from an online prioritization exercise involving 168 managers from 22 different European countries. In general, the topics of major concern for forest managers coincided with the ones that are at the heart of most research projects. They covered important issues related to the management of mixed forests and the role of mixtures for the stability of forest functioning and management identified and selected by a range of European forest managers during an extensive participatory process. The set of 10 questions were the highest ranked questions from an online prioritization exercise involving 168 managers from 22 different European countries. In general, the topics of major concern for forest managers coincided with the ones that are at the heart of most research projects. They covered important issues related to the management of mixed forests and the role of mixtures for the stability of forests faced with environmental changes and the provision of ecosystem services to society. Our analysis showed that the current scientific knowledge about these questions was rather variable and particularly low for those related to the management of mixed forests over time and the associated costs. We also found that whereas most research projects have sought to evaluate whether mixed forests are more stable or provide more goods and services than monocultures, there is still little information on the underlying mechanisms and trade-offs behind these effects. Similarly, we identified a lack of knowledge on the spatio-temporal scales at which the effects of mixtures on the resistance and adaptability to environmental changes are operating. Our analysis may help researchers to identify what knowledge needs to be better transferred and to better design future research initiatives meeting practitioner's concerns.

Keywords: Species mixtures, Review, Forest management and functioning, Participatory process, Research challenges, Ecosystem services, Forest stability

1. Introduction

In recent years, the study of mixed forests has been the focus of increasing research efforts, in particular the consequences of admixing tree species for the productivity and stability of forest systems. This has generated a substantial amount of new knowledge (e.g. Pretzsch et al., 2013; Vilà et al., 2013; Morin et al., 2014; Tobner et al., 2016; Liang et al., 2016; van der Plas et al., 2016; among others), and the consolidation of important scientific initiatives and networks (Baeten et al., 2013; Bravo-Oviedo et al., 2014; Verheyen et al., 2016). From the research perspective, the recent advances in the understanding of mixed forests functioning are of unquestionable value, but the extent to which this information is responding to practitioners' concerns remains unknown.

We addressed this issue via a collaborative work in the context of the EuMIXFOR research network (Bravo-Oviedo et al., 2014) in which researchers from 30 different European countries participated. The study was divided into three steps. First, we conducted a Pan-European survey with the objective of identifying key questions related to mixtures that, from the perspective of forest managers, still require further research attention. Second, we ranked these questions by relevance according to the views of an independent set of European practitioners obtained via an online prioritization exercise. Finally, we evaluated current scientific knowledge for the highest ranked questions and we identified future research challenges in relation to them. The ultimate aim of our work was to reduce the commonly reported gap between knowledge generated from research and that required by forest

managers (see Petrokofsky et al., 2010). In that respect, we expect our analysis will provide both (i) information to the research community on the priority knowledge needs of forest practitioners and (ii) brief reviews of the current state of knowledge regarding the topics of their concern. Finally, we expect that the identification of research challenges (based on the questions received from the practitioners) may help researchers to contextualise and design future research initiatives and may also facilitate the translation of new knowledge into practical outcomes.

2. Collection and prioritization of research questions by forest managers

2.1. Collection of questions

Each representative of the individual European countries that participated in the *EUMIXFOR* network contacted forest managers from that country who had expertise in the management of mixed-forests in either public or private ownership. We asked the managers to provide a list of the 5–10 key questions about mixtures for which they would like more information from the research community (preferably in the form of an interrogative sentence). Fifty-three forest managers from 15 countries responded to this request providing 289 questions (Fig. 1). The set of questions from each country was added sequentially to the pool of questions. The sets of questions identified by the last countries added to the list did not raise additional topics, suggesting that the main questions had already been covered and that adding new

STEP 1: COLLECTION OF QUESTIONS

53 managers from 15 countries were asked to provide a list of the 5–10 key questions about mixtures



289 questions

STEP 2: CLASSIFICATION

A research multidisciplinary team classified, merged and rephrased the questions

30 questions



STEP 3: PRIORITIZATION

168 managers from 22 countries participated to a questionnaire to identify the most important research questions

10 questions

Fig. 1. Schematic representation of the participatory process conducted with European forest managers for the selection of the 10 questions used to structure the review. The countries colored in green corresponded to the ones that contributed to step 1 (above) and step 3 (below).

countries would not increase the number of questions to be retained.

A multidisciplinary group of six experienced forest researchers (LC, CC, ML, BM, QP and KV) within the network classified each question into eleven broad themes (e.g. timber production, species interactions...) during a one-day workshop. Questions within each theme were then combined (when overlapping) and rephrased (if they were unclearly formulated or related to a very specific type of mixture) by this group of researchers. During this process, the only questions discarded were those that did not relate to mixtures. The process concluded with the formulation of 30 questions covering most of the replies originally received (Table S1).

2.2. Prioritization process

These 30 questions related to mixed forests were then ranked through an online prioritization survey conducted in 22 countries throughout Europe (Fig. 1). We contacted an independent sample of 168 forestry professionals (i.e. between 5 and 15 forest managers per country), working in different organisations (public institutions, private forests, forest associations) and with a professional interest in the management of mixtures. We presented the 30 questions (translated into their national language) to each of the 168 respondents that participated in the exercise, and we used the best-worst scaling (BWS) method to rank them according to the preferences of each individual.

The BWS method (Finn and Louviere, 1992; Louviere et al., 2013) is a discrete choice task in which each respondent is asked repeatedly to choose the most important and the least important item from among randomly selected subsets of the original set of items, in this case of 4 out of the 30 questions. BWS forces respondents to discriminate among the presented alternatives, thus preventing some of the problems associated with other ranking methodologies, such as anchoring bias, i.e. the tendency of respondents to consistently use the middle points or one of the end points when using rating scales (Flynn et al., 2007; Rudd and Lawton, 2013). The prioritization exercise was conducted using an internet-based survey platform (SurveyGizmo, Boulder, CO, USA).

The values ascribed to the different questions ranged from nearly 63 for the highest ranked to about 39 for the lowest ranked questions (Table S1). A feature of the exercise was that a number of questions given an upper to middle ranking (e.g. ranks 8–18) received quite similar scores. In order to constrain the length of the review section that follows, we took an arbitrary decision to limit detailed discussion to the ten most highly ranked questions. Similar procedures of constraining results of participatory processes to the ten highest questions have been used in other studies (e.g. Petrovsky et al., 2010).

3. Revision of the current state of knowledge in relation to forest managers' questions

We synthesize below the current state of knowledge in relation to the ten highest ranked questions selected by forest managers. The questions were categorized into three broad groups as they refer to the relation between mixed forests and (i) stability, (ii) the provision of ecosystem services, and (iii) management. The questions within each group were addressed in the order we considered the most appropriate to facilitate the flow of writing and reading. In the sections below, the number in brackets next to each question shows its rank that resulted from the prioritization process (see Table S1).

3.1. Stability

- Which mixtures of species provide the best resistance and best resilience to climate change and natural disturbances? (#1)
- Are mixed forests more resistant and resilient to climate change and natural disturbances? (#2)

In recent years, the question of whether mixed forests are better able

to cope with environmental change than monocultures has been a focus of attention (see for example the reviews by Thompson et al., 2009; Bauhus and Schmerbeck, 2010 or Scherer-Lorenzen, 2014). The concepts of resilience and resistance have been addressed and defined in many different ways (Brand, 2009). Here, we follow the approach of Hodgson et al. (2015) and we consider resilience to encompass both resistance and recovery; with the first being the capacity of the system to absorb an exogenous disturbance and the second its capacity to come back to an equilibrium after being disturbed (see also Oliver et al., 2015). Forest resilience can be approached at the level of periodic stresses (e.g. drought episodes) or of disturbances (e.g. windstorms, fires) (see Trumbore et al., 2015). In the case of most European forests, there is a large consensus that the impacts of both types of stressor are expected to increase with climate change (Seidl et al., 2011). The response of forests to periodic stresses relates to the concept of ecosystem stability, a concept that has been largely investigated in grassland ecosystems, where diversity helps to maintain the productivity of ecosystems subject to climate variations (Tilman et al., 2006; Isbell et al., 2015). The diversity-stability relationship in forest ecosystems is less clear (Thompson et al., 2009), although some comprehensive studies such as the ones by Morin et al. (2014) and Jucker et al. (2014) also reported more stable productivity of mixed-forests over time. Such stabilizing effects might be mediated by a reduction of the competition among species for growing resources (i.e. functional complementarity (Loreau and Hector, 2001)), asynchronic species-intrinsic responses to environmental fluctuations (Morin et al., 2014) or by temporal shifts in species interactions (i.e. temporal complementarity) (del Rio et al., 2017).

Forest resistance to biotic factors, such as insect herbivores or fungal pathogens, increases in mixed-forests which in general present lower pest abundance and experience lesser damage than monocultures (see meta-analysis by Jactel et al., 2005 or Haas et al., 2011). These findings are explained by different mechanisms such as reduced host tree density and accessibility ("associational resistance hypothesis", Barbosa et al., 2009), or by an increased presence of predators and parasitoids in more diverse forests (Guyot et al., 2016). However, reduced damage by insect herbivores in mixed forests is not observed consistently (see for example Vehviläinen et al., 2006; Schuldt et al., 2010; Haase et al., 2015) and the same occurs with fungal disease incidence (Nguyen et al., 2016). In some cases, reversed patterns (i.e. higher damage in mixed forests) have been reported when damages are triggered by generalist herbivores ("associational susceptibility hypothesis", Barbosa et al., 2009). Some authors have concluded that biotic damages are in many cases more related to the specific composition of the forests (or the type of herbivore) than to species richness *per se* (see meta-analysis by Vehviläinen et al., 2007 or Jactel and Brockerhoff, 2007). Similar conclusions derive from the few existing studies investigating the impact of mammal herbivores in mixed stands (Vehviläinen and Koricheva, 2006, Metslaid et al., 2013).

Similarly to biotic damages, the role of tree diversity in the capacity of forests to resist severe abiotic disturbances (such as catastrophic windstorms or wildfires) is unclear and appears to be more dependent on structure and species combinations than on diversity (Dhôte, 2005; Grossiord et al., 2014; Pereira et al., 2014; Forrester et al., 2016; Metz et al., 2016). In contrast, tree diversity is generally considered to enhance the capacity of forests to recover from disturbances although this has been tested in field studies since it requires long-term monitoring and adequate information about the state of the forest prior to the disturbances. The higher resilience of mixtures to severe disturbances might be mediated by the higher diversity and higher redundancy of traits relevant to tree response to environmental changes (e.g. resprouting capacity, seed bank longevity) that these stands may present (Yachi and Loreau, 1999; Laliberté et al., 2010; Puettmann, 2011; Sánchez-Pinillos et al., 2016).

From a management perspective, promoting the coexistence of species belonging to different functional groups and/or with different

strategies to face disturbances (to increase the probability of recovery processes) seems a good starting point (Sánchez-Pinillos et al., 2016). This mostly translates into trying to maintain the inherent complexity of forests, i.e. to develop (wherever possible) within- and among-stand heterogeneity in ecosystem structure, composition, and to accept variability in space and time as an inherent attribute to enhance forests' natural capacity to adapt and self-organize in response to gradual or abrupt environmental changes (Lloret et al., 2007; Puettmann et al., 2009; Messier et al., 2013).

3.2. Provision of ecosystem services

Forest ecosystem services are the range of benefits people obtain from forests. They include provisioning, regulating, cultural and supporting services (MEA, 2005) and arise from ecosystem functions provided by organisms (Scherer-Lorenzen, 2014). Understanding the influence of biodiversity on ecosystem services requires analysing (i) the ecological processes that produce the ecosystem functions and (ii) the economic and sociological processes that value these functions into services that eventually provide human well-being (Butterfield et al., 2016).

Among forest ecosystem services, wood production has been the most studied service, but other services such as soil protection, plant and animal diversity, carbon sequestration and their relationship to tree diversity are currently being investigated in forest biomes.

- How do mixed forests affect the quantity and quality of wood production? (#5)

Several meta-analyses and reviews accounting for confounding factors such as site, species pool and stand characteristics, have shown an overall positive Diversity-Productivity Relationship (DPR) in forest ecosystems at stand/plot scale (typically < 0.1 ha) (Paquette and Messier, 2011; Bauhus and Schmerbeck, 2010; Zhang et al., 2012; Liang et al., 2016). On average, stand production is higher in a mixture compared to expectation based on the mean production in pure stands of the component species, yet some individual monocultures may still be more productive than the most productive mixtures.

To value the wood volume produced and evaluate the socio-economic impact of tree diversity, it is necessary to sort the wood volume produced into wood quality classes, which correspond to particular classes of use and may be assigned a specific economic value. In a recent review, Pretzsch and Rais (2016) reported that the effects of tree diversity on wood quality were balanced and ambiguous, since tree morphology, structure and wood quality are strongly affected by stand structural heterogeneity, which is generally higher in a mixed than in a pure stand (see also Zeller et al., 2017).

- Are mixed-forests more efficient in using resources (light, water, nutrients) than pure ones? (#10)

Positive DPRs are related to selection (when changes in the relative yields of species in a mixture are non-randomly related to their yields in monoculture; Loreau and Hector, (2001)) and complementarity resulting from (i) competitive reduction (when competition is reduced in mixtures compared to pure stands) or (ii) facilitation (when a species improves the functioning of another species) (Vandermeer, 1989). Complementarity arises from inter-specific differences in physiology, phenology or morphology or from intra-specific differences that result from inter-specific interactions, and is affected by stand structure (Richards et al., 2010; Forrester and Bauhus, 2016). There is important variability among DPRs, even for a given species pool. The Monteith primary production model may be used as a framework to explain how the slope of the DPR changes along spatial or temporal gradients in resource availability or climatic conditions (Forrester and Bauhus, 2016). Complementarity is predicted to increase as the availability of a

given resource declines (or as climatic conditions become harsher) if interactions among associated species result in an improvement of the availability, uptake or use-efficiency of that resource (or if interactions improve the climatic condition). Functional differences among admixed species appear to be a key condition for overyielding to occur (Zhang et al., 2012), but the net effect of these functional differences on overyielding depends on how they can reduce climate constraints/increase availability of limiting resources on a particular site.

- Do mixed-forests provide more ecosystem services than monocultures? (#9)

3.2.1. Carbon sequestration

The effects of tree species diversity on C sequestration may be assessed by considering (i) the biologically-mediated processes that drive the rates of C gain and loss and the size and longevity of C stocks, and (ii) the processes that determine the associated social and economic values (Diaz et al., 2009a; Diaz et al., 2009b). While the contribution of tree diversity to the net C uptake in aboveground tree components may be derived from DPRs, its impacts on belowground C storage, including roots and soils, remain much less documented (Hulvey et al., 2013). Because trade-offs at the individual tree species level prevent the maximizing of C sequestration across multiple C pools (e.g. root vs shoot biomass; Hulvey et al., 2013), maximizing forest C sequestration is expected to be achieved by using selected combinations of species traits. The complex effects of tree species diversity and identity on C storage are well illustrated when analysing soil C stocks. Dawud et al. (2016) observed a limited influence of tree species diversity and identity on the overall C soil storage (0–40 cm), but contrasting effects on the distribution of C within the soil profile. Diversity tended to increase C in deeper layers; by contrast, the effect of diversity on the forest floor C stock was inconsistent, in agreement with Handa et al. (2014) who clearly showed that the functional diversity of both decomposers and leaf litter, not simply litter species richness, promotes C and N cycling. As opposed to diversity, species identity tended to influence C storage in the upper forest floor layers. If confirmed by other studies, tree species diversity would therefore mainly benefit the longevity of C stocks through its effects on C storage in the deeper soil layers.

3.2.2. Plant and animal diversity

Canopy trees represent only a small part of forest biodiversity. The impacts of tree diversity on plant, animal and fungal diversity are complex. On one hand, mixed forests can be more productive, they also present higher structural heterogeneity which may provide more diverse above- and belowground microhabitats than monocultures, and may therefore host a greater number of organisms (De Deyn et al., 2004). On the other hand, neutral or negative effects of tree diversity may be observed in mixed forest where a dilution of each individual tree species may eliminate organisms that are dependent on particular tree species (Ampoorter et al., 2014; Tederloo et al., 2016). In a literature review, Cavard et al. (2011) examined existing empirical evidence that tree mixtures promote the diversity of understory plants, songbird, soil fauna, and ectomycorrhiza in northern forests. They found no evidence of the existence of organisms uniquely associated with mixtures, species richness simply reflecting, at best, the accumulation of organisms associated with each canopy tree species. They also reported that tree diversity improves the diversity of understory plants (but see Barbier et al., 2008), avian and ectomycorrhizal communities (see also Bibby et al., 1989). Although many studies found positive effects of mixtures on earthworm or microarthropod diversity (see Korboulewsky et al., 2016), no general trend emerged on the relationship between mixed forests and soil fauna diversity.

3.2.3. Provision of multiple ecosystem services

Many studies have focused on the relationships between tree diversity and individual forest ecosystem functions, but very few studies

have examined the impacts of tree diversity on ecosystem services, and even fewer studies have analysed multiple functions and services.

Multifunctional forest management requires that multiple ecosystem functions and services are simultaneously sustained. Several studies, mainly from grassland experiments, demonstrated that the level of biodiversity needed to maintain multiple functions was greater than the levels needed to maximize each individual function (Hector and Bagchi, 2007; Lefcheck et al., 2015); considering multiple locations and long time series in a changing environment further increases the needed level of biodiversity to provide multiple functions (Isbell et al., 2011).

The degree of multifunctionality of a forest can be determined by the number of ecosystem functions exceeding a predefined threshold value (Byrnes et al., 2014). Using such an approach, van der Plas et al. (2016) showed that multifunctionality increased with species richness for moderate levels of functioning, while it decreased when high function levels are desired. One may therefore conclude that the simultaneous maximisation of all functions at a stand level is not achievable as a result of trade-off between functions.

- Which mixture of species (or functional groups) should be used to optimize specific or combined management targets (e.g. productivity, biodiversity, stability...)? (#4)
- Which positive and negative effects on different ecosystem functions (e.g. productivity, litter decomposition, stem quality) can occur when mixing particular species? (#6)

Although many ecosystem functions are on average positively associated with canopy tree diversity (Nadrowski et al., 2010), there is often a considerable scattering around the mean, and for a given diversity level, the outcome of the interactions may be either positive, neutral or even negative, depending on the identities of the associated species (Scherer-Lorenzen, 2014). Moreover, even when similar species are combined, the outcome still depends on the set of current environmental conditions, including resource availability and climate constraints, as reported above for DPRs. From the manager's perspective, this means that effective tree species selection has to consider not only the functional differences between the investigated species for those traits involved in the function of interest, but also how functional diversity is expected to translate into positive effects given the environmental conditions at hand. While approaches using functional diversity metrics (Laliberté and Legendre, 2010; Mouchet et al., 2010) and empirical frameworks relating complementarity to resource availability and climate (Forrester and Bauhus, 2016) may assist optimal species selection, process-based models, such as those developed for growth (Forrester and Tang, 2016), appear quite promising as they combine the most relevant mechanisms and their interactions.

Regarding the optimization of combined management targets, van der Plas et al. (2016) showed that the relationship between multifunctionality and tree species richness described above was driven by the 'Jack-of-all-trades' effect, with only minor effects of either 'complementarity' or 'selection'. This means that whenever species effects on different functions are not perfectly correlated, the functioning of a multi-species mixture equals the biomass-weighted average of the function levels of monocultures of its component species.

For some functions, however, the relationship with tree species diversity remains much less documented or general patterns have not been discerned (Nadrowski et al., 2010). This is the case, among others, for those functions and processes that are more strongly affected by site conditions such as belowground processes and biogeochemical cycling (Scherer-Lorenzen, 2014). In addition to the identity effects discussed above, the possible context dependency of the Diversity Ecosystem functions Relationships (DERs) could also explain the lack of net diversity effects when encompassing a range of sites, contrasting DERs slopes between sites being driven by environmental factors.

3.3. Management

- What silvicultural treatments should be applied to maintain the desired species throughout the entire stand rotation? (#3)

The silvicultural treatments applied to any mixture should reflect the management objectives chosen for the forest while respecting edaphic factors and species composition and characteristics. A useful framework for evaluating the potential effectiveness of silvicultural interventions at different phases of stand development is provided by a model of stand dynamics (Oliver and Larson, 1996) which separates stand development into four stages: stand initiation, stem exclusion, understorey reinitiation and old-growth (note that the last stage is rare in many managed forests). The creation of mixtures is best achieved in the first and third stages, whereas in the second stage thinning is used to ensure the survival of an existing mixture. However, at all stages, careful tending can be essential to ensure that the balance of a desired mixture is maintained.

During the stand initiation stage, acceptance of natural regeneration of a range of species that are suited to the site is often the best and most cost-effective way of developing a mixed stand. This approach can be combined with planting so that the regeneration forms the matrix between planted groups of a desired species (Saha et al., 2013), or can be favoured to create a two storied stand (Frivold and Groven, 1996; Stanturf et al., 2014). Two-storied mixed stands can also be created by deliberately underplanting fast growing pioneer tree species with slower growing and shade tolerant broadleaves or conifers (Pommerening and Murphy, 2004; Kelty, 2006; Paquette and Messier, 2013). Planting of mixtures is an option on nutrient poor soils where a more nutrient demanding species is mixed with one adapted to such sites, as is the case for the pine/spruce mixtures reported from the British Isles (Gabriel et al., 2005; Mason and Connolly, 2014) and Poland (Bielak et al., 2014) or where a nitrogen fixing species is mixed with another valuable timber species such as walnut (*Juglans regia* L.) or *Eucalyptus* spp. (Clark et al., 2008; Forrester et al., 2011; Radosevich et al., 2006).

Once the trees have closed canopy (stem exclusion), a period of intense inter-tree competition begins which can be mediated by the selective removal of individual trees or species (a.k.a 'thinning'). Where species are of compatible growth rates and shade tolerance, there is little need to adjust thinning strategies from practice in pure stands. The challenge occurs where the competition from one species can disadvantage the growth of a favoured species, as occurs with aspen (*Populus tremuloides* Michx.) and white spruce (*Picea glauca* (Moench) Voss) in boreal mixedwoods (Filipescu and Comeau, 2007). In such instances, thinning will need to favour stems of a more vulnerable but desirable species by removing immediate competitors. Other examples include mixtures of oak and more shade tolerant tree species (such as beech) where thinning is mandatory to prevent the latter outcompeting the more valuable oak (Hein and Dhôte, 2006; Johnson et al., 2009).

As the trees age, the canopy either begins to open up naturally or small gaps are created through thinning deliberate thinning regime or by the final harvest. As a result, the increased light on the forest floor allows tree seedlings of a range of species to become established ('understorey reinitiation'). With control of ungulate browsing and careful tending, over time such seedlings (planted or naturally regenerated) can be promoted into the upper canopy layers and can be used to help convert a regular structure to an irregular one (Mosandl and Kleinert, 1998; Knoke and Plusczyk 2001; Nyland, 2003; O'Hara, 2014). This process can be used as a means of converting pure planted stands to mixed irregular forests, as in the conversion of Norway spruce to mixed conifer-broadleaved stands in some regions of central and western Europe (Spiecker et al., 2004; Ammer et al., 2008) or in restoring natural forest types after larch afforestation in northern China (Mason and Zhu, 2014). The development and formation of these mixed stands can be fostered by a range of irregular silvicultural systems (Matthews,

Table 1

List of the 10 high-ranked questions resulting from the participatory process with European managers. For each question the current level of scientific knowledge is evaluated as follows: + (hardly any research results available), ++ (individual case-studies available), +++ (integrative studies, reviews or meta-analyses available). Some key references and research needs are also provided.

Rank-position	Question	Current knowledge	Some key references	Research needs
#1	Which mixtures of species provide the best resistance and best resilience to climate change and natural disturbances?	+	Pretzsch et al. (2013); Sánchez-Pinillos et al. (2016)	Role of different components of biodiversity (species richness, functional diversity) and organizational levels (e.g. trophic levels)
#2	Are mixed forests more resistant and resilient to climate change and natural disturbances?	+++	Jactel et al. (2005); Neuner et al. (2015)	Disturbance interactions and cascading effects; cross-scale approaches
#3	What silvicultural treatments should be applied to maintain the desired species throughout the entire stand rotation?	+	Pommerening and Murphy, (2004)	Establishment and analysis of long-term research plots
#4	Which mixture of species (or functional groups) should be used to optimize specific or combined management targets (e.g. productivity, biodiversity, stability...)?	++	Scherer Lorenzen, (2014); van der Plas et al. (2016)	Translation of individual and combined ecosystem functions into ecosystem services; long-term research plots
#5	How do mixed forests affect the quantity and quality of wood production?	+++*	Vilà et al. (2013); Pretzsch and Rais, (2016)	Factors behind transgressive overyielding of mixtures; effects of the mixture composition and stand structure
#6	Which positive and negative effects on different ecosystem functions (e.g. productivity, litter decomposition, stem quality) can occur when mixing particular species?	++	Nadrowski et al. (2010)	Impact of mixtures on belowground processes and biogeochemical cycles; interactions between belowground and aboveground responses; context dependency
#7	How does the expected balance of benefits and costs compare between pure and mixed stands?	++	Knoke et al. (2008); Neuner et al. (2015)	of the relationship between diversity and ecosystem functions Integrated economic analyses with inclusion of uncertainty and risk (timber price fluctuations, disturbance occurrence)
#8	Do mixtures allow more flexibility and provide more options to adapt to changing management objectives than monocultures?	+	—	Analyses of documented case studies; operational-scale demonstrations
#9	Do mixed-forests provide more ecosystem services than monocultures?	++	Gamfeldt et al. (2013)	Impact of mixtures on belowground processes and biogeochemical cycles
#10	Are mixed-forests more efficient in using resources (light, water, nutrients) than pure ones?	+++	Forrester, (2014); Forrester and Bauhus, (2016)	Development of process-based models for mixed stands;

* Refers to the level of knowledge on the relation between mixtures and the quantity of wood production. The existing knowledge in relation to the effects of mixtures on wood quality is much lower (+).

1991) involving combinations of tree species of different functional traits. While the general principles of the transformation process outlined above are well understood, their formulation into silvicultural guidelines for the management of particular species combinations in specific site conditions is often lacking. In part, this major knowledge gap reflects the historic emphasis given to experimentation with single species stands which means that the complexities of successfully manipulating species mixtures over time are poorly described and little known.

- Do mixtures allow more flexibility and provide more options to adapt to changing management objectives than monocultures? (#8)

Conceptually, the presence of more than one species in a maturing stand should give forest managers greater flexibility to adapt to changing objectives and to harvest different products at different stages of a stand's development (Nichols et al., 2006). However, it is difficult to find cases where this theoretical benefit has actually been realised or where there has been a comparison with pure stands. One example occurred in the UK in the 1960s when policy for public forests changed from developing a strategic supply of timber for the market to maximizing the return on investment. As a result, a silvicultural regime for management of nursing mixtures of conifers and broadleaves in lowland Britain (Kerr et al., 1992) was changed from gradually removing the conifers to favour the broadleaves to one of eliminating the broadleaves to favour the faster growing conifers. The occurrence of aspen and white spruce in either two or single storey mixtures in boreal Canada is another example where the combination can allow managers to harvest either species for different products depending on market conditions and demand (Comeau et al., 2005).

- How does the expected balance of benefits and costs compare between pure and mixed stands? (#7)

For forest managers, any evaluation of benefits and costs from mixtures is heavily dependent on financial returns from wood production rather than involving consideration of wider aspects such as the relative delivery of ecosystem services (Quine et al., 2013). Establishment costs can heavily influence the potential profitability of mixtures. Saha et al. (2013), for example, showed that group plantings of oak in broadleaved regeneration were cheaper to establish and maintain than conventional pure oak planting in an analysis carried out in young (10–26 years old) forest stands of central and southern Germany. Comparisons of the relative returns from pure and mixed stands depend upon the anticipated yields from the two types of stands, and a situation where a high yielding species is mixed with a less productive one often results in lower total yield and a reduction in theoretical profits (Knoke et al., 2008). However, if the probability of risks from disturbances (biotic or abiotic), which are generally higher for pure stands, are calculated (e.g. Neuner et al., 2015) it can be shown that the mixed stand has a higher outturn, especially for a risk averse investor/owner and where longer rotations are incurred (Roessiger et al., 2013). In addition, a yield stimulus of 10%, depending on product and rotation length, can offset any increased costs associated with planting and managing mixed-species stands (Nichols et al., 2006). For example, if proper allowance is made for any positive yield improvement from growing species in mixture, then the financial performance of the mixture is better than that of the pure stand, as in two-storied mixtures of birch (*Betula pendula* Roth. and *Betula pubescens* Ehrh.) and Norway spruce (*Picea abies* (L.) Karsten) in Scandinavia (Valkonen and Valsta, 2001). However, such results can be influenced by stand structure since the financial outturn from single storied mixed stands of the same species was lower in the mixture than in the pure stand (Fahlvik et al., 2011). These results highlight how evaluation of the relative balance of the financial return from mixtures can be context dependent, influenced by factors such as forest type and owner objectives (Felton et al., 2016).

4. General discussion and future research directions

We summarise above the current state of knowledge in relation to the ten highest ranked questions related to mixed-forest management and functioning that are of major concern from the view of European forest managers. Our exercise could be conceived as a discussion between research suppliers and users: we consider that it has delivered results of high interest for both groups. The questions for which forest managers showed the most concern related to the capacity of mixed forests to respond to the effects of climate change and/or to the occurrence of natural disturbances. This could be explained by the recognized uncertainty of, and unpredictability associated with, these events and to the fact that they are not “controllable” by the implementation of any management strategy or action. Interestingly, these topics have been at the centre of many research initiatives (see Table 1). There is a general agreement in the scientific literature that mixtures are more resilient to natural disturbances than monocultures and that they present more options for adaptation to climate change. However, some of these positive aspects seem to be more related to the specific composition of the mixture than to tree diversity *per se* (see for example Metz et al., 2016), and additional efforts should be undertaken to assess which combination of species or functional groups needs to be promoted to tackle potential negative effects of predicted (or unexpected) environmental changes. Indeed, we share the view of Jactel et al. (2016) that further research efforts in this topic might be devoted to the understanding of potential trade-offs between species and communities with regards to the resistance and recovery to different disturbances and environmental changes. Improving our understanding of the spatio-temporal scales at which the effects of mixtures on the resistance and adaptability to change are operating might also be considered in future research projects (Table 1).

In contrast to the analysis of the underlying mechanisms behind the diversity – stability relationship, which has received substantial attention from the research community, we have poor information on how to manage tree mixtures over time and the cost (and benefits) behind these systems. Accordingly, we were able to provide very few evidence-based responses to the questions raised by the managers in relation to this area. Once the scarce published literature on this topic was reviewed, we observed that there is a critical lack of long-term research plots that explore and illustrate the silviculture of mixed forests in different forest types (Table 1). Such plots are necessary to validate the results of more theoretical studies as well as to support practice and the development of guidelines for the management of mixed forests. They could also be used to address whether browsing pressure from ungulates prevents or not the establishment of new mixed-species plantations (e.g. Bergquist et al., 2009). We also recognized there are almost no documented case studies which provide operational evidence of the greater management flexibility presumed to be provided by mixed forests, and very few integrated economic analysis showing the effects of a greater use of mixtures on the provision of ecosystem services within the forestry-wood chain. Such analyses may need to take proper account of uncertainty and risk and to provide costs and revenues which are relevant to managers' needs (Table 1).

Our survey also revealed the interest of forest managers in receiving research evidence about the widespread view that mixed forests provide more ecosystem functions and services than monocultures (five out of the ten highest ranked questions on mixed forests were related to this topic). The analysis we conducted confirmed this statement. Knowledge about tree species diversity effects on forest functioning has increased considerably in recent years resulting in general principles that could be translated into guidelines to be used by forest practitioners (Forrester and Bauhus, 2016).

However, we still lack integrated assessments of the role of the various components of biodiversity (e.g. species richness, species composition, community evenness, functional diversity, phylogenetic diversity) as well as of the organizational levels (trophic levels, taxa/

organisms, ...) on the provision of ecosystem functions (and in particular to those related to belowground processes and biogeochemical cycles) (Table 1). Indeed, we are still far from understanding how individual and combined ecosystem functions translate into ecosystem services. We also detected the need for further understanding of the biodiversity-ecosystem function relationship at all relevant temporal and spatial scales for management issues, while still accounting for confounding factors. Studies dealing with the response of forest ecosystem functions to biodiversity are often restricted to the stand scale (but see Chisholm et al., 2013), and to a very limited fraction of the stand cycle and tree lifespan. Lastly, we consider that additional efforts need to be devoted to the development of process-based models to help forest managers define best tree species combinations to optimize the supply of targeted services (while keeping the others at relatively high levels) (Table 1). For operational use, these models should provide managers with accurate information on product outturn, wood properties and timber value.

In conclusion, the results of our analysis show a general agreement between forest managers' concerns and the topics that are at the heart of most research projects dealing with mixed-forests. However, we have detected substantial differences in the amount of available knowledge relating to the various questions provided by the managers. Whereas most research projects have sought to evaluate whether mixed forests provide more goods and services than monocultures and are more stable when faced with environmental change (i.e. the effects of mixing, questions #2, #5), there is still little information about the underlying mechanisms and trade-offs behind these effects (although these questions are currently at the heart of a number of research initiatives (Verheyen et al., 2016)). Finally, our results stress the critical need of generating additional knowledge to provide forest managers with evidence-based silvicultural guidelines allowing the establishment and maintenance of mixtures over time under different environmental conditions.

Acknowledgements

This article is based upon work from COST Action FP1206, EuMIXFOR, supported by COST (European Cooperation in Science and Technology). The authors want to specially thank the large number of forest managers that were involved in the participatory process and kindly responded to the surveys. The first author also thanks the Spanish Ministry of Economy and Competitiveness for funding the project AGL2015-70425-R which partly supported his contribution to this study. MM acknowledges support by grant IUT21-4 of the Estonian Ministry of Education and Research and DS support by project III 43007 (2011–2017) from Ministry of Education, Science and Technological Development of Republic of Serbia.

References

Ammer, C., Bickel, E., Kölling, C., 2008. Converting Norway spruce stands with beech—a review of arguments and techniques. *Aust. J. For. Sci.* 125, 3–26.

Ampoorter, E., Baeten, L., Koricheva, J., Vanhellemont, M., Verheyen, K., 2014. Do diverse overstoreys induce diverse understoreys? Lessons learnt from an experimental-observational platform in Finland. *For. Ecol. Manage.* 318, 206–215.

Baeten, L., Verheyen, K., Wirth, C., Bruehlheide, H., Bussotti, F., Finér, L., Jaroszewicz, B., Selvi, F., Valladares, F., Allan, E., Ampoorter, E., Auge, H., Avacariev, D., Barbaro, L., Barnoaiea, I., Bastias, C.C., Bauhus, J., Beinhoff, C., Benavides, R., Benneter, A., Berger, S., Berthold, F., Boberg, J., Bonal, D., Brüggemann, W., Carnol, M., Castagnayrol, B., Charbonnier, Y., Checko, E., Coomes, D., Coppi, A., Dalmaris, E., Danila, G., Dawud, S.M., de Vries, W., De Wandeler, H., Deconchat, M., Domisch, T., Duduman, G., Fischer, M., Fotelli, M., Gessler, A., Gimeno, T.E., Granier, A., Grossiord, C., Guyot, V., Hantsch, L., Hättenschwiler, S., Hector, A., Hermý, M., Holland, V., Jactel, H., Joly, F.-X., Jucker, T., Kolb, S., Koricheva, J., Lexer, M.J.,

Liebigesell, M., Milligan, H., Müller, S., Muys, B., Nguyen, D., Nichiforel, L., Pollastrini, M., Proulx, R., Rabasa, S., Radoglou, K., Ratcliffe, S., Raulund-Rasmussen, K., Seiferling, I., Stenlid, J., Vesterdal, L., von Wilpert, K., Zavala, M.A., Zielinski, D., Scherer-Lorenzen, M., 2013. A novel comparative research platform designed to determine the functional significance of tree species diversity in European forests. *Perspect. Plant Ecol. Evol. Syst.* 15, 281–291.

Barbier, S., Gosselin, F., Balandier, P., 2008. Influence of tree species on understory vegetation diversity and mechanisms involved – a critical review for temperate and boreal forests. *For. Ecol. Manage.* 254 (1), 1–15.

Barbosa, P., Hines, J., Kaplan, I., Martinson, H., Szczepaniec, A., Szendrei, Z., 2009. Associational resistance and associational susceptibility: having right or wrong neighbors. *Annu. Rev. Ecol. Evol. Syst.* 40, 1–20.

Bauhus, J., Schmerbeck, J., 2010. Silvicultural options to enhance and use forest plantation biodiversity. In: Bauhus, J., van der Meer, P., Kanninen, M. (Eds.), *Ecosystem Goods and Services from Plantation Forests*. Earthscan, London, pp. 96–139.

Bergquist, J., Löf, M., Örlander, G., 2009. Effects of roe deer browsing and site preparation on performance of planted broadleaved and conifer seedlings when using temporary fences. *Scand. J. For. Res.* 24, 308–317.

Bibby, C.J., Aston, N., Bellamy, P.E., 1989. Effects of broadleaved trees on birds of upland conifer plantations in North Wales. *Biol. Conserv.* 49, 17–29.

Bielak, K., Dudzinska, M., Pretzsch, H., 2014. Mixed stands of Scots pine (*Pinus sylvestris* L.) and Norway spruce [*Picea abies* (L.) Karst] can be more productive than monocultures. Evidence from over 100 years of observation of long-term experiments. *For. Syst.* 23, 573–589.

Brand, F., 2009. Critical natural capital revisited: Ecological resilience and sustainable development. *Ecol. Econ.* 68, 605–612.

Bravo-Oviedo, A., Barreiro, S., Strelcova, K., Pretzsch, H., 2014. EuMIXFOR Introduction: integrating scientific knowledge in sustainable forest management of mixed forests. *For. Syst.* 23, 515–517.

Butterfield, B.J., Camhi, A.L., Rubin, R.L., Schwalm, C.R., 2016. Tradeoffs and compatibilities among ecosystem services: biological, physical and economic drivers of multifunctionality. In: Woodward, G., Bohan, D.A. (Eds.), *Advances in Ecological Research*. Academic Press, Oxford, pp. 207–243.

Byrnes, J.E.K., Gamfeldt, L., Isbell, F., Lefcheck, J.S., Griffin, J.N., Hector, A., Cardinale, B.J., Hooper, D.U., Dee, L.E., Duffy, J.E., 2014. Investigating the relationship between biodiversity and ecosystem multifunctionality: challenges and solutions. *Methods Ecol. Evol.* 5, 111–124.

Cavard, X., Macdonald, S.E., Bergeron, Y., Chen, H.Y.H., 2011. Importance of mixed-woods for biodiversity conservation: evidence for understory plants, songbirds, soil fauna, and ectomycorrhizae in northern forests. *Environ. Rev.* 19, 142–161.

Chisholm, R.A., Muller-Landau, H.C., Rahman, K.A., Bebb, D.P., Bin, Y., Bohlman, S.A., Bourg, N.A., Brinks, J., Bunyavechewin, S., Butt, N., Cao, H., Cao, M., Cárdenas, D., Chang, L.-W., Chiang, J.-M., Chuyong, G., Condit, R., Dattaraja, H.S., Davies, S., Duque, A., Fletcher, C., Gunatilleke, N., Gunatilleke, S., Hao, Z., Harrison, R.D., Howe, R., Hsieh, C.-F., Hubbell, S.P., Itoh, A., Kenfack, D., Kiratiprayoon, S., Larson, A.J., Lian, J., Lin, D., Liu, H., Lutz, J.A., Ma, K., Malhi, Y., McMahon, S., McShea, W., Meegaskumbura, M., Mohd, R.S., Morecroft, M.D., Nyctch, C.J., Oliveira, A., Parker, G.G., Pulla, S., Punchi-Manage, R., Romero, S.H., Sang, W., Schurman, J., Su, S.-H., Sukumar, R., Sun, I.-F., Suresh, H.S., Tan, S., Thomas, D., Thomas, S., Thompson, J., Valencia, R., Wolf, A., Yap, S., Ye, W., Yuan, Z., Zimmerman, J.K., 2013. Scale-dependent relationships between tree species richness and ecosystem function in forests. *J. Ecol.* 101, 1214–1224.

Clark, J.R., Hemery, G.E., Savill, P.S., 2008. Early growth and form of common walnut (*Juglans regia* L.) in mixture with tree and shrub nurse species in southern England. *Forestry* 81, 631–644.

Comeau, P.G., Kabzems, R., McClarnon, J., Heineman, J.L., 2005. Implications of selected approaches for regenerating and managing western boreal mixedwoods. *For. Chron.* 81, 559–574.

Dawud, S.M., Raulund-Rasmussen, K., Domisch, T., Finér, L., Jaroszewicz, B., Vesterdal, L., 2016. Is tree species diversity or species identity the more important driver of soil carbon stocks, C/N ratio, and pH? *Ecosystems* 19, 645–660.

De Deyn, G.B., Raaijmakers, C.E., van Ruijven, J., Berendse, F., van der Putten, W.H., 2004. Plant species identity and diversity effects on different trophic levels of nematodes in the soil food web. *Oikos* 106, 576–586.

Dhôte, J.-F., 2005. Implications of forest diversity in resistance to strong winds. In: Scherer Lorenzen, M., Korner, C., Schulze, E.-D. (Eds.), *Forest Diversity and Function Temperate and Boreal Systems*. Ecological Studies. Springer Vol. 176. Berlin, Germany, pp. 291–307.

Díaz, S., Hector, A., Wardle, D.A., 2009a. Biodiversity in forest carbon sequestration initiatives: not just a side benefit. *Curr. Opin. Environ. Sustain.* 1, 55–60.

Díaz, S., Wardle, D.A., Hector, A., 2009b. Incorporating biodiversity in climate change mitigation initiatives. In: Naeem, S., Bunker, D.E., Hector, A., Loreau, M., Perrings, C. (Eds.), *Biodiversity, Ecosystem Functioning, and Human Wellbeing: An Ecological and Economic Perspective*. Oxford University Press, UK, pp. 149–166.

Fahvik, N., Agestam, E., Ekö, P.M., Linden, M., 2011. Development of single-storied mixtures of Norway spruce and birch in Southern Sweden. *Scand. J. For. Res.* 26, 36–45.

Felton, A., Nilsson, U., Sonesson, J., Felton, A.M., Roberge, J.-M., Ranius, T., Ahlström, M., Bergh, J., Björkman, C., Boberg, J., Drössler, L., Fahvik, N., Gong, P., Holmström, E., Kesikitalo, E.C.H., Klapwijk, M.J., Laudon, H., Lundmark, T., Niklasson, M., Nordin, A., Pettersson, M., Stenlid, J., Sténs, A., Wallertz, K., 2016. Replacing monocultures with mixed-species stands: ecosystem service implications of two production forest alternatives in Sweden. *Ambio* 45, 124–139.

Filipescu, C.N., Comeau, P.G., 2007. Aspen competition affects light and white spruce growth across several boreal sites in western Canada. *Can. J. For. Res.* 37, 1701–1713.

- Finn, A., Louviere, J.J., 1992. Determining the appropriate response to evidence of public concern: the case of food safety. *J. Pub. Pol. Mark* 11, 12–25.
- Flynn, T.N., Louviere, J.J., Peters, T.J., Coast, J., 2007. Best–worst scaling: what it can do for health care research and how to do it. *J. Health Econ.* 26, 171–189.
- Forrester, D.I., Vancly, J.K., Forrester, R.I., 2011. The balance between facilitation and competition in mixtures of *Eucalyptus* and *Acacia* changes as stands develop. *Oecologia* 166, 265–272.
- Forrester, D.I., 2014. The spatial and temporal dynamics of species interactions in mixed-species forests: from patterns to process. *For. Ecol. Manage.* 312, 282–292.
- Forrester, D.I., Tang, X., 2016. Analysing the spatial and temporal dynamics of species interactions in mixed-species forests and the effects of stand density using the 3-PG model. *Ecol. Mod.* 319, 233–254.
- Forrester, D.I., Bauhus, J., 2016. A review of processes behind diversity-productivity relationships in forests. *Curr. For. Rep.* 2, 45–61.
- Forrester, D.I., Bonal, D., Dawud, S., Gessler, A., Granier, G., Pollastrini, M., Grossiord, C., 2016. Drought responses by individual tree species are not often correlated with tree species diversity in European forests. *J. Appl. Ecol.* 53, 1725–1734.
- Frivold, L.H., Groven, R., 1996. Yield and management of mixed stands of spruce, birch and aspen. *Nor. J. Agr. Sci. supp.* 24, 1–21.
- Gabriel, K., Blair, I., Mason, W.L., 2005. Growing broadleaved trees on the North York Moors: results after nearly 50 years. *Q. J. For.* 99, 21–30.
- Gamfeldt, L., Snäll, T., Bagchi, R., Jonsson, M., Gustafsson, L., Kjellander, P., Ruiz-Jaen, M.C., Fröberg, M., Stendahl, J., Philipson, C.D., Mikusiński, G., Andersson, E., Westerlund, B., Andrén, H., Moberg, F., Moen, J., Bengtsson, J., 2013. Higher levels of multiple ecosystem services are found in forests with more tree species. *Nat. Comm.* 4, 1340.
- Grossiord, C., Granier, A., Ratcliffe, S., Bouriaud, O., Bruehlheide, H., Cheeko, E., Forrester, D.I., Dawud, S.M., Finér, L., Pollastrini, M., Scherer-Lorenzen, M., Valladares, F., Bonal, D., Gessler, A., 2014. Tree diversity does not always improve resistance of forest ecosystems to drought. *Proc. Natl. Acad. Sci.* 111, 14812–14815.
- Guyot, V., Castagneyrol, B., Vialatte, A., Deconchat, M., Jactel, H., 2016. Tree diversity reduces pest damage in mature forests across Europe. *Biol. Lett.* 12, 20151037.
- Haase, J., Castagneyrol, B., Cornelissen, J.H.C., Ghazoul, J., Kattge, J., Koricheva, J., Scherer-Lorenzen, M., Morath, S., Jactel, H., 2015. Contrasting effects of tree diversity on young tree growth and resistance to insect herbivores across three biodiversity experiments. *Oikos* 124, 1674–1685.
- Haas, S.E., Hooten, M.B., Rizzo, D.M., Meentemeyer, R.K., 2011. Forest species diversity reduces disease risk in a generalist plant pathogen invasion. *Ecol. Lett.* 14, 1108–1116.
- Handa, I.T., Aerts, R., Berendse, F., Berg, M.P., Bruder, A., Butenschoten, O., Chauvet, E., Gessner, M.O., Jabiol, J., Makkonen, M., McKie, B.G., Malmqvist, B., Peeters, E.T.H.M., Scheu, S., Schmid, B., van Ruijven, J., Vos, V.C.A., Hättenschwiler, S., 2014. Consequences of biodiversity loss for litter decomposition across biomes. *Nature* 509, 218–221.
- Hector, A., Bagchi, R., 2007. Biodiversity and ecosystem multifunctionality. *Nature* 448, 188–190.
- Hein, S., Dhôte, J.F., 2006. Effect of species composition, stand density and site index on the basal area increment of oak trees (*Quercus* sp.) in mixed stands with beech (*Fagus sylvatica* L.) in northern France. *Ann. For. Sci.* 63, 457–467.
- Hodgson, D., McDonald, J.L., Hosken, D.J., 2015. What do you mean, resilient? *Trends Ecol. Evol.* 30, 503–506.
- Hulvey, K.B., Hobbs, R.J., Standish, R.J., Lindenmayer, D.B., Lach, L., Perring, M.P., 2013. Benefits of tree mixes in carbon plantings. *Nat. Clim. Chang.* 3, 869–874.
- Isbell, F., Calcagno, V., Hector, A., Connolly, J., Harpole, W.S., Reich, P.B., Scherer-Lorenzen, M., Schmid, B., Tilman, D., van Ruijven, J., Weigelt, A., Wilsey, B.J., Zavaleta, E.S., Loreau, M., 2011. High plant diversity is needed to maintain ecosystem services. *Nature* 477, 199–202.
- Isbell, F., Craven, D., Connolly, J., Loreau, M., Schmid, B., Beierkuhnlein, C., Bezemer, T.M., Bonin, C., Bruehlheide, H., de Luca, E., Ebeling, A., Griffin, J.N., Guo, Q., Hauri, Y., Hector, A., Jentsch, A., Kreyling, J., Lanta, V., Manning, P., Meyer, S.T., Mori, A.S., Naeem, S., Niklaus, P.A., Polley, H.W., Reich, P.B., Roscher, C., Seabloom, E.W., Smith, M.D., Thakur, M.P., Tilman, D., Tracy, B.F., van der Putten, W.H., van Ruijven, J., Weigelt, A., Weisser, W.W., Wilsey, B., Eisenhauer, N., 2015. Biodiversity increases the resistance of ecosystem productivity to climate extremes. *Nature* 526, 574–547.
- Jactel, H., Brockerhoff, E.G., Duelli, P., 2005. A test of the biodiversity-stability theory: meta-analysis of tree species diversity effects in insect pest infestations, and re-examination of responsible factors. In: Scherer Lorenzen, M., Korner, C., Schulze, E.-D., (Eds.), *Forest Diversity and Function: Temperate and Boreal Systems. Ecological Studies*, vol. 176, Springer, Berlin, Germany, pp. 235–261.
- Jactel, H., Brockerhoff, E.G., 2007. Tree diversity reduces herbivory by forest insects. *Ecol. Lett.* 10, 835–848.
- Jactel, H., Boberg, J., Bonal, D., Castagneyrol, B., Gardiner, B., Gonzalez-Olabarria, J.R., Koricheva, J., Meurisse, N., Brockerhoff, E.G., 2016. Tree diversity-forest resistance relationships. In: *Integrating Scientific Knowledge in Mixed Forests (Book of abstracts of the EuMIXFOR Final Conference)*, 5–7 October, Prague, Czech Republic.
- Johnson, P.S., Shifley, S.R., Rogers, R., 2009. *The Ecology and Silviculture of Oaks*. CABI Publishing, New York.
- Jucker, T., Bouriaud, O., Avacaritei, D., Coomes, D.A., 2014. Stabilizing effects of diversity on aboveground wood production in forest ecosystems: linking patterns and processes. *Ecol. Lett.* 17, 1560–1569.
- Kelty, M.J., 2006. The role of species mixtures in plantation forestry. *For. Ecol. Manage.* 233, 195–204.
- Kerr, G., Nixon, C.J., Matthews, R.W., 1992. Silviculture and yield of mixed-species stands: the UK experience. In: Cannell, M.G.R., Malcolm, D.C., Robertson, P.A. (Eds.), *The Ecology of Mixed-Species Stands of Trees*. Blackwell, Oxford, pp. 35–51.
- Knoke, T., Plusczyk, N., 2001. On economic consequences of transformation of a spruce (*Picea abies* (L.) Karst.) dominated stand from regular into irregular age structure. *For. Ecol. Manage.* 151, 163–179.
- Knoke, T., Ammer, C., Stimm, B., Mosandl, R., 2008. Admixing broadleaved to coniferous tree species: A review on yield, ecological stability and economics. *Eur. J. For. Res.* 127, 89–101.
- Korboulewsky, N., Perez, G., Chauvat, M., 2016. How tree diversity affects soil fauna diversity: a review. *Soil Biol. Biochem.* 94, 94–106.
- Laliberté, E., Legendre, P., 2010. A distance-based framework for measuring functional diversity from multiple traits. *Ecology* 91, 299–305.
- Laliberté, E., Wells, J.A., Declerck, F., Metcalfe, D.J., Catterall, C.P., Queiroz, C., Aubin, I., Bonser, S.P., Ding, Y., Fraterrigo, J.M., McNamara, S., Morgan, J.W., SánchezMerlos, D., Vesk, P.A., Mayfield, M.M., 2010. Land-use intensification reduces functional redundancy and response diversity in plant communities. *Ecol. Lett.* 13, 76–86.
- Lefcheck, J.S., Byrnes, J.E.K., Isbell, F., Gamfeldt, L., Griffin, J.N., Eisenhauer, N., Hensel, M.J.S., Hector, A., Cardinale, B.J., Duffy, J.E., 2015. Biodiversity enhances ecosystem multifunctionality across trophic levels and habitats. *Nat. Comm.* 6, 6936.
- Liang, J., Crowther, T.W., Picard, N., Wiser, S., Zhou, M., Alberti, G., Schulze, E.-D., McGuire, A.D., Bozzato, F., Pretzsch, H., de-Miguel, S., Paquette, A., Hérault, B., Scherer-Lorenzen, M., Barrett, C.B., Glick, H.B., Hengeveld, G.M., Nabuurs, G.-J., Pfautsch, S., Viana, H., Vibrans, A.C., Ammer, C., Schall, P., Verbyla, D., Tchebakova, N., Fischer, M., Watson, J.V., Chen, H.Y.H., Lei, X., Schelhaas, M.-J., Lu, H., Gianelle, D., Parfenova, E.I., Salas, C., Lee, E., Lee, B., Kim, H.S., Bruehlheide, H., Coomes, D.A., Pionto, D., Sunderland, T., Schmid, B., Gourlet-Fleury, S., Sonké, B., Tavani, R., Zhu, J., Brandt, S., Vayreda, J., Kitahara, F., Searle, E.B., Neldner, V.J., Ngugi, M.R., Baraloto, C., Frizzera, L., Balazy, R., Oleksyn, J., Zawila-Niedzwiecki, T., Bouriaud, O., Bussotti, F., Finér, L., Jaroszewicz, B., Jucker, T., Valladares, F., Jagodzinski, A.M., Peri, P.L., Gonmadje, C., Marthy, W., O'Brien, T., Martin, E.H., Marshall, A., Rovero, F., Bitariho, R., Niklaus, P.A., Alvarez-Loayza, P., Chamuya, N., Valencia, R., Mortier, F., Wortel, V., Engone-Obiang, N.L., Ferreira, L.V., Odeke, D.E., Vasquez, R.M., Reich, P.B., 2016. Positive biodiversity-productivity relationship predominant in global forests. *Science* 354 (6309), 196.
- Lloret, F., Lobo, A., Estevan, H., Maisongrande, P., Vayreda, J., Terradas, J., 2007. Woody plant richness and NDVI response to drought events in Catalanian (northeastern Spain) forests. *Ecology* 88, 2270–2279.
- Loreau, M., Hector, A., 2001. Partitioning selection and complementarity in biodiversity experiments. *Nature* 412, 72–76.
- Louviere, J., Lings, I., Islam, T., Gudergan, S., Flynn, T., 2013. An introduction to the application of (case 1) best-worst scaling in marketing research. *Int. J. Res. Mark.* 30, 292–303.
- Mason, W.L., Zhu, J.J., 2014. Silviculture of planted forests managed for multi-functional objectives: lessons from Chinese and British experiences. In: Fenning, T. (Ed.), *Challenges and Opportunities for the World's Forests in the 21st Century*. Springer 81, New York, pp. 37–54.
- Mason, W.L., Connolly, T., 2014. Mixtures with spruce species can be more productive than monocultures: evidence from the Gisburn experiment in Britain. *Forestry* 87, 209–217.
- Matthews, J.D., 1991. *Silvicultural Systems*. Clarendon Press, Oxford.
- Messier, C., Puettmann, K.J., Coates, K.D., 2013. Managing forests as complex adaptive systems – building resilience to the challenge of global change. Routledge, New York.
- Metslaid, M., Palli, T., Randveer, T., Sims, A., Jögi, K., Stanturf, J.A., 2013. The condition of Scots pine stands in Lahemaa National Park, Estonia 25 years after browsing by moose (*Alces alces*). *Boreal Environ. Res.* 18, 25–34.
- Metz, J., Annighöfer, P., Schall, P., Zimmermann, J., Kahl, T., Schulze, E.D., Ammer, C., 2016. Site-adapted admixed tree species reduce drought susceptibility of mature European beech. *Glob. Chang. Biol.* 22, 903–920.
- Millennium Ecosystem Assessment, 2005. *Ecosystems and Human Well-being: Synthesis*. Island Press, Washington, DC.
- Morin, X., Fahse, L., De Mazancourt, C., Scherer-Lorenzen, M., Bugmann, H., 2014. Diversity enhances the temporal stability of forest productivity in time because of stronger asynchrony in species dynamics. *Ecol. Lett.* 17, 1526–1535.
- Mosandl, R., Kleinert, A., 1998. Development of oaks (*Quercus petraea* (Matt.) Liebl.) emerged from bird-dispersed seeds under old-growth pine (*Pinus sylvestris* L.) stands. *For. Ecol. Manage.* 106, 35–44.
- Mouchet, M.A., Villeger, S., Mason, N.W., Mouillot, D., 2010. Functional diversity measures: an overview of their redundancy and their ability to discriminate community assembly rules. *Funct. Ecol.* 24, 867–876.
- Nadrowski, K., Wirth, C., Scherer-Lorenzen, M., 2010. Is forest diversity driving ecosystem function and service? *Curr. Opin. Environ. Sustain.* 2, 75–79.
- Neuner, S., Albrecht, A., Cullmann, D., Engels, F., Griess, V.C., Hahn, W.A., Hanewinkel, M., Härtl, F., Kölling, C., Staupendahl, K., Knoke, T., 2015. Survival of Norway spruce remains higher in mixed stands under a dryer and warmer climate. *Global Ch. Biol.* 21, 935–946.
- Nguyen, D., Castagneyrol, B., Bruehlheide, H., Bussotti, F., Guyot, V., Jactel, H., Jaroszewicz, B., Valladares, F., Stenlid, J., Boberg, J., 2016. Fungal disease incidence along tree diversity gradients depends on mixture in European forests. *Ecol. Evol.* 6, 2426–2438.
- Nichols, J.D., Bristow, M., Vancly, J.K., 2006. Mixed-species plantations: prospects and challenges. *For. Ecol. Manage.* 233, 383–390.
- Nyland, R.D., 2003. Even- to uneven-aged: the challenges of conversion. *For. Ecol. Manage.* 172, 291–300.
- O'Hara, K., 2014. *Multitaged Silviculture*. Oxford University Press, United Kingdom, Managing for Complex Forest Stand Structures.
- Oliver, C.D., Larson, B.C., 1996. *Forest Stand Dynamics*. John Wiley and Sons, New York.
- Oliver, T.M., Heard, M.S., Isaac, N.J.B., Roy, D.B., Procter, D., Eigenbrod, F., Freckleton, R., Hector, A., Orme, C.D.L., Petchey, O.L., Proença, V., Raffaelli, D., Suttle, K.B.,

- Mace, G.M., Martín-López, B., Woodcock, B.A., Bullock, J.M., 2015. Biodiversity and resilience of ecosystem functions. *Trends Ecol. Evol.* 30, 673–684.
- Paquette, A., Messier, C., 2011. The effect of biodiversity on tree productivity: from temperate to boreal forests. *Glob. Ecol. Biogeogr.* 20, 170–180.
- Paquette, A., Messier, C., 2013. Managing Tree Plantations as Complex Adaptive Systems. In: Messier, C., Puettmann, K.J., Coates, K.D. (Eds.), *Managing forests as complex adaptive systems: Building Resilience to the Challenge of Global Change*, Routledge, Earthscan, New York, pp. 299–326.
- van der Plas, F., Manning, P., Allan, E., Scherer-Lorenzen, M., Verheyen, K., Wirth, C., Zavala, M.A., Hector, A., Ampoorter, E., Baeten, L., Barbaro, L., Bauhus, J., Benavides, R., Benneter, A., Berthold, F., Bonal, D., Bouriaud, O., Bruelheide, H., Bussotti, F., Carnol, M., Castagneyrol, B., Charbonnier, Y., Coomes, D., Coppi, A., Bastias, C.C., Muhie Dawud, S., De Wandeler, H., Domisch, T., Finer, L., Gessler, A., Granier, A., Grossiord, C., Guyot, V., Hattenschwiler, S., Jactel, H., Jaroszewicz, B., Joly, F.-X., Jucker, T., Koricheva, J., Milligan, H., Muller, S., Muys, B., Nguyen, D., Pollastrini, M., Raulund-Rasmussen, K., Selvi, F., Stenlid, J., Valladares, F., Vesterdal, L., Zielinski, D., Fischer, M., 2016. Jack-of-all-trades effects drive biodiversity–ecosystem multifunctionality relationships in European forests. *Nat. Comm.* 7 (11109).
- Pereira, M.G., Aranha, J., Amraoui, M., 2014. Land cover fire proneness in Europe. *For. Syst.* 23, 598–610.
- Petrokofsky, G., Brown, N.D., Hemery, G.E., Woodward, S., Wilson, E., Weatherall, A., Stokes, V., Smithers, R.J., Sangster, M., Russell, K., Pullin, A.S., Price, C., Morecroft, M., Malins, M., Lawrence, A., Kirby, K.J., Godbold, D., Charman, E., Boshier, D., Bosbeer, S., Arnold, J.E.M., 2010. A participatory process for identifying and prioritizing policy-relevant research questions in natural resource management: a case study from the UK forestry sector. *Forestry* 83, 357–367.
- Pommerening, A., Murphy, S.T., 2004. A review of the history, definitions and methods of continuous cover forestry with special attention to afforestation and restocking. *Forestry* 77, 27–44.
- Pretzsch, H., Schütze, G., Uhl, E., 2013. Resistance of European tree species to drought stress in mixed versus pure forests: evidence of stress release by inter-specific facilitation. *Plant Biol.* 15, 483–495.
- Pretzsch, H., Rais, A., 2016. Wood quality in complex forests versus even-aged monocultures: review and perspectives. *Wood Sci. Technol.* 50, 845–880.
- Puettmann, K.J., Coates, K.D., Messier, C., 2009. *A Critique of Silviculture: Managing for Complexity*. Island Press, Washington, D.C.
- Puettmann, K.J., 2011. Silvicultural challenges and options in the context of global change: “Simple” fixes and opportunities for new management approaches. *J. For.* 109, 321–331.
- Quine, C.P., Bailey, S.A., Watts, K., 2013. Sustainable forest management in a time of ecosystem services frameworks: common ground and consequences. *J. Appl. Ecol.* 50, 863–867.
- Radosevich, S., Hibbs, D., Ghersa, C., 2006. Effects of species mixtures on growth and stand development of Douglas-fir and red alder. *Can. J. For. Res.* 36, 768–782.
- Richards, A.E., Forrester, D.I., Bauhus, J., Scherer-Lorenzen, M., 2010. The influence of mixed tree plantations on the nutrition of individual species: a review. *Tree Physiol.* 30, 1192–1208.
- del Río, M., Pretzsch, H., Ruiz-Peinado, R., Ampoorter, E., Annighöfer, P., Barbeito, I., Bielak, K., Brazaitis, G., Coll, L., Drössler, L., Fabrika, M., Forrester, D., Heym, M., Hurt, V., Kurylyak, V., Löf, M., Lombardi, F., Makrickiene, E., Matovic, B., Mohren, F., Motta, R., van Ouden, J., Pach, M., Ponette, Q., Schütze, G., Skrzyszewski, J., Sramek, V., Sterba, H., Stojanovic, D., Svoboda, M., Zlatanov, T., Bravo-Oviedo, A., 2017. Species interactions increase the temporal stability of community productivity in *Pinus sylvestris*-*Fagus sylvatica* mixtures across Europe. *J. Ecol.* <http://dx.doi.org/10.1111/1365-2745.12727>.
- Roessiger, J., Griess, V.C., Härtl, F., Clasen, C., Knoke, T., 2013. How economic performance of a stand increases due to decreased failure risk associated with the admixing of species. *Ecol. Model.* 255, 58–69.
- Rudd, M.A., Lawton, R.N., 2013. Scientists’ prioritization of global coastal research questions. *Mar. Policy* 39, 101–111.
- Saha, S., Kuehne, C., Bauhus, J., 2013. Tree species richness and stand productivity in low-density cluster plantings with oaks (*Quercus robur* L. and *Q. petraea* (Mattuschka) Liebl.). *Forests* 4, 650–665.
- Sánchez-Pinillos, M., Coll, L., De Cáceres, M., Ameztegui, A., 2016. Assessing the persistence capacity of communities facing natural disturbances on the basis of species response traits. *Ecol. Ind.* 66, 76–85.
- Scherer-Lorenzen, M., 2014. The functional role of biodiversity in the context of global change. In: Coomes, D.A., Burslem, D.F.R.P., Simonson, W.D. (Eds.), *Forests and Global Change*. Cambridge University Press, UK, pp. 195–237.
- Schuldt, A., Baruffol, M., Böhnke, M., Bruelheide, H., Härdtle, W., Lang, A.C., Nadrowski, K., Von Oheimb, G., Voigt, W., Zhou, H., Assmann, T., 2010. Tree diversity promotes insect herbivory in subtropical forests of south-east China. *J. Ecol.* 98, 917–926.
- Seidl, R., Schelhaas, M.-J., Lexer, M.J., 2011. Unraveling the drivers of intensifying forest disturbance regimes in Europe. *Glob. Chang. Biol.* 17, 2842–2852.
- Spiecker, H., Hansen, J., Klimo, E., Skovgaard, J.P., Sterba, H., Teuffel, K.V., 2004. Norway Spruce Conversion-options and Consequences. *Eur. For. Inst. Res. Rep.* 18, 1–269.
- Stanturf, J.A., Palik, B.J., Dumroese, R.K., 2014. Contemporary forest restoration: a review emphasizing function. *For. Ecol. Manage.* 331, 292–323.
- Tedersoo, L., Bahram, M., Cajthaml, T., Pölme, S., Hiesalu, I., Anslan, S., Harend, H., Buegger, F., Pritsch, K., Koricheva, J., Abarenkov, K., 2016. Tree diversity and species identity effects on soil fungi, protists and animals are context dependent. *Int. Soc. Microb. Ecol. J.* 10, 346–362.
- Thompson, I., Mackey, B., McNulty, S., Mosseler, A., 2009. *Forest Resilience, Biodiversity, and Climate Change*. Secretariat of the Convention on Biological Diversity, Montreal. Technical Series no. 43.
- Tilman, D., Reich, P.B., Knops, J.M.H., 2006. Biodiversity and ecosystem stability in a decade-long grassland experiment. *Nature* 441, 629–632.
- Tobner, C.M., Paquette, A., Gravel, D., Reich, P.B., Williams, L., Messier, C., 2016. Functional identity drives overyielding in early tree communities. *Ecol. Lett.* 19, 638–647.
- Trumbore, S., Brando, P., Hartmann, H., 2015. Forest health and global change. *Science* 349, 814–818.
- Valkonen, S., Valsta, L., 2001. Productivity and economics of mixed two-storied spruce and birch stands in Southern Finland simulated with empirical models. *For. Ecol. Manage.* 140, 133–149.
- Vandermeer, J., 1989. *The Ecology of Intercropping*. Cambridge University Press.
- Vehviläinen, H., Koricheva, J., Ruohomäki, K., Johansson, T., Valkonen, S., 2006. Effects of tree stand species composition on insect herbivory of silver birch in boreal forests. *Basic Appl. Ecol.* 7, 1–11.
- Vehviläinen, H., Koricheva, J., 2006. Moose and vole browsing patterns in experimentally assembled pure and mixed forest stands. *Ecography* 29, 497–506.
- Vehviläinen, H., Koricheva, J., Ruohomäki, K., 2007. Tree species diversity influences herbivore abundance and damage: meta-analysis of long-term forest experiments. *Oecologia* 152, 287–298.
- Verheyen, K., Vanhellemont, M., Auge, H., Baeten, L., Baraloto, C., Barsoum, N., Bilodeau-Gauthier, S., Bruelheide, H., Castagneyrol, B., Godbold, D., Haase, J., Hector, A., Jactel, H., Koricheva, J., Loreau, M., Mereu, S., Messier, C., Muys, B., Nolet, P., Paquette, A., Parker, J., Perring, M., Ponette, Q., Potvin, C., Reich, P., Smith, A., Weih, M., Scherer-Lorenzen, M., 2016. Contributions of a global network of tree diversity experiments to sustainable forest plantations. *Ambio* 45, 29–41.
- Vilà, M., Carrillo-Gavilán, A., Vayreda, J., Bugmann, H., Fridman, F., Grodzki, W., Haase, J., Kunstler, G., Schelhaas, A., Trasobares, A., 2013. Disentangling biodiversity and climatic determinants of wood production. *Plos One* 8, e53530.
- Yachi, S., Loreau, M., 1999. Biodiversity and ecosystem productivity in a fluctuating environment: the insurance hypothesis. *Proc. Natl. Acad. Sci.* 96, 1463–1468.
- Zeller, L., Ammer, C., Annighöfer, P., Biber, P., Marshall, J., Schütze, G., del Río Gaztelurrutia, M., Pretzsch, H., 2017. Tree ring wood density of Scots pine and European beech lower in mixed-species stands compared with monocultures. *For. Ecol. Manage.* 400, 363–374.
- Zhang, Y., Chen, H.Y.H., Reich, P.B., 2012. Forest productivity increases with evenness, species richness and trait variation: a global meta-analysis. *J. Ecol.* 100, 742–749.

Supplementary information

Table S1. List of 30 questions ordered by their rank value (expressed on a 0–100 scale) after the prioritization exercise

	Question formulation	Rank-value
#1	Which mixtures of species provide the best resistance and best resilience to climate change and natural disturbances?	62,98
#2	Are mixed forests more resistant and resilient to climate change and natural disturbances?	58,88
#3	What silvicultural treatments should be applied to maintain the desired species throughout the entire stand rotation?	58,39
#4	Which mixture of species (or functional groups) should be used to optimize specific or combined management targets (e.g. productivity, biodiversity, stability...)?	58,21
#5	How do mixed forests affect the quantity and quality of wood production?	57,46
#6	Which positive and negative effects on different ecosystem functions (e.g. productivity, litter decomposition, stem quality) can occur when mixing particular species?	55,84
#7	How does the expected balance of benefits and costs compare between pure and mixed stands?	55,24
#8	Do mixtures allow more flexibility and provide more options to adapt to changing management objectives than monocultures?	53,84
#9	Do mixed-forests provide more ecosystem services than monocultures?	53,68
#10	Are mixed-forests more efficient in using resources (light, water, nutrients) than pure ones?	52,76
#11	How do effects of mixed-forest effects on productivity and resilience change along stand developmental stages?	52,49
#12	What stand structural and spatial patterns should be favoured to maintain mixtures of species with contrasting shade tolerance?	52,42
#13	What are the best options to convert monocultures to mixtures?	52,30
#14	How can the ecological impacts and benefits of mixed-forests be quantified?	52,01
#15	Are there adequate models to predict the growth and management of complex mixed stands?	51,51
#16	Do intimate mixtures provide more (or different) benefits compared to patch or landscape scale mixtures?	50,57
#17	What are the most appropriate harvesting systems for use in mixed forests?	50,53
#18	Are there some site conditions that are more suitable for promoting tree species mixtures and for obtaining any associated benefits?	49,59
#19	What are the impacts of tree-species mixtures on soils at the stand and ecosystem levels?	48,20
#20	How much does biodiversity increase if we increase the number of tree species in the stand?	47,77
#21	How do we establish mixed species stands as part of afforestation programmes?	46,77
#22	Is there a minimum threshold in terms of species proportion required to induce a mixing effect at the stand level?	45,88
#23	Is it possible to predict the impacts of mixing on ecosystem- / stand- level properties based on the traits of the associated tree species?	45,54
#24	How do effects of mixed-forest on productivity and resilience change along abiotic gradients?	45,06
#25	Do we need improved sampling methods for use in inventories in mixed forests?	41,92
#26	Is there a desirable (optimal) balance to be achieved between the amount of pure and mixed stands at the landscape or regional level?	41,62
#27	What are the impacts of mixing on individual tree functioning (water status, nutrition)?	41,15
#28	Can any mixed species stands be sustained without management?	40,54
#29	Can the fragmentation characteristic of private forests lead to practical problems when managing mixed forests?	40,13
#30	What are the impacts of mixtures of provenances within tree species on ecosystem functioning (compared to those expected from mixtures of tree species)?	38,89