



Article

Organic Fertilizer Sources Distinctively Modulate Productivity, Quality, Mineral Composition, and Soil Enzyme Activity of Greenhouse Lettuce Grown in Degraded Soil

Mariateresa Cardarelli ¹, Antonio El Chami ¹, Paola Iovieno ², Youssef Rouphael ³, Paolo Bonini ⁴ and Giuseppe Colla ^{1,*}

¹ Department of Agriculture and Forest Sciences, University of Tuscia, 01100 Viterbo, Italy

² CREA—Consiglio per la Ricerca in Agricoltura e l'Analisi dell'Economia Agraria, Centro di Ricerca Orticoltura e Florovivaismo, 84098 Pontecagnano Faiano, Italy

³ Department of Agricultural Sciences, University of Naples Federico II, 80055 Portici, Italy

⁴ oloBion S.L., 08028 Barcelona, Spain

* Correspondence: giucolla@unitus.it

Abstract: Intensive greenhouse vegetable production is often associated with a decline of crop productivity due to the increase of soil salinity and/or a reduction of biological fertility. The aim of the current work was to assess the effects of three organic fertilizers on morpho-physiological and agronomic traits of greenhouse lettuce as well as soil enzyme activity under poor soil quality conditions. The tested organic fertilizers (poultry manure, vinasse-based fertilizer, and insect's frass fertilizer) were applied pre-planting at the same equivalent nitrogen (N) rate (90 kg N ha⁻¹). Laboratory incubation assay results showed that vinasse-based fertilizer was the most suitable fertilizer in supplying the mineral N in the short term. All fertilizers increased shoot fresh and dry weight compared to unfertilized control with a more pronounced effect (+75%) with vinasse-based fertilizer and insect's frass. Insect frass reduced by 27% the leaf nitrate concentration in comparison with the other treatments. The toxic heavy metal Pb was 46% lower in all organically fertilized lettuce leaves. Soil enzymatic activities of acid phosphatase, alkaline phosphatase, arylsulfatase (ArS), N-acetyl-β-D-glucosaminidase (NAGase), dehydrogenase, and total hydrolase (THA) were enhanced by poultry manure and insect's frass in comparison with unfertilized control while vinasse-based fertilizer increased ArS, NAGase, and THA. Taken together, our data demonstrate that the application of organic fertilizers especially vinasse-based fertilizer and insect's frass during intensive crop production is a suitable approach for mitigating the negative impact of soil salinity, enhancing soil biological fertility, and improving agronomic performance of greenhouse lettuce.

Keywords: organic fertilizers; *Lactuca sativa* L.; physiological traits; produce quality; enzyme activity; sustainable horticulture



Citation: Cardarelli, M.; El Chami, A.; Iovieno, P.; Rouphael, Y.; Bonini, P.; Colla, G. Organic Fertilizer Sources Distinctively Modulate Productivity, Quality, Mineral Composition, and Soil Enzyme Activity of Greenhouse Lettuce Grown in Degraded Soil.

Agronomy **2023**, *13*, 194. <https://doi.org/10.3390/agronomy13010194>

Academic Editor: Claudio Ciavatta

Received: 8 December 2022

Revised: 2 January 2023

Accepted: 4 January 2023

Published: 7 January 2023



Copyright: © 2023 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

1. Introduction

Finding efficient means to optimize crop growth and quality without squandering priceless natural resources is one of the major issues affecting the entire food supply chain today. The preservation of significant non-renewable resources, such as soil, should come after efforts to reduce chemical (e.g., synthetic) inputs and maintain production quality. Soil quality and fertility are being negatively impacted by anthropogenic activity and climate change, which also result in issues with organic matter loss, rising salinity from chemical abuse, and drought from insufficient precipitation [1]. Traditional and intensive cultivation methods can increase the salt content in the soil, and crops cultivated directly in the soil can be negatively affected by low content of organic matter. Soil salinization and loss of organic matter are the main causes of soil degradation in vegetable crop production. The public widely criticized the greenhouse industry for using unnatural production techniques and for severely polluting

the environment with the residues of fertilizers, pesticides, and other agrochemicals used [2]. Therefore, a sustainable intensification of crop production which aims maintaining a good soil fertility is very important for assuring long-term crop productivity.

The European Union is making serious efforts to provide tools to reduce agriculture's environmental impact, starting with legislation [3]. The circular economy foundations encourage the reuse and valorization of by-products from the agricultural and livestock industry to create suitable products for cultivations, such as organic fertilizers. The rise of organic agriculture is one of the main efforts to make this sector more sustainable, increasingly avoiding synthetic fertilizers and pesticides [4]. Several studies have shown that using organic fertilizers results in an improvement of soil physical properties, soil enzyme activity, soil organic carbon, and an increase in nutrients which leads to enhance microbial development, thus increasing soil fertility and quality [5–8]. Soil enzymes and soil microorganisms play an active role in influencing soil fertility due to their participation in the cycle of nutrients, such as carbon and nitrogen which are required for plant growth, but they are also sensitive biological indicators for soil quality evaluation in addition to sensitively reflecting minute changes in soil environment [9,10].

Sustainable organic fertilization relies on understanding how the quality of organic fertilizer affects the synchrony of nitrogen (N) supply and crop N demand which is critical for optimal agronomic N management, environmental protection, and crop productivity, especially in short crop cycles [11,12]. Some authors investigated the rate of soil respiration and the dynamics of the mineral N in various soil/organic fertilizer mixtures in a laboratory incubation experiment [13]. When compared to urea, organic fertilizers significantly increased soil respiration and slowed the release of mineral N in the soil. Moreover, the dynamic of mineral N released from organic fertilizers varied significantly during the incubation period (12 weeks) with a fast mineral N release at 25 °C in the organic fertilizers containing feather meal, vinasse, guano, and poultry manure. Organic fertilizers with a fast release of N are necessary to meet the N requirements of short life cycle crops such as lettuce. Lettuce is a fast-growing crop which requires a continuous and sufficient supply of nutrients during all growing cycles. Moreover, lettuce is sensitive to soil salinity [14] causing growth depression and physiological disorder known by lettuce tip burn that is due to a problem of calcium uptake and transportation in leaves during periods of rapid growth [15]. In addition, as a leafy crop, lettuce is very susceptible to nitrate accumulation in its leaves due to the oversupply with nitrogen fertilizers [16,17]. Nitrate accumulation in edible plant tissue has a significant impact on human health; when it is converted to nitrite in the body, it can result in respiratory dysfunction called methemoglobinemia, which is harmful to children [18]. Furthermore, it might also oversee the formation of nitrosamines, which are known to cause cancer [19].

Researchers have focused on investigating the effects of different organic fertilizers on lettuce growth, yield, leaf nitrate content, and soil fertility, while limited studies have considered the agronomic effect of organic fertilizers under low fertile soils resulting from intensive greenhouse horticulture. Moreover, limited studies have considered newly insect frass fertilizers derived from the digestate of black soldier fly larvae. A group of authors evaluated the potential agronomic value of black soldier fly larvae frass (BSFF) as organic fertilizer in greenhouse lettuce production under pot conditions in comparison with mineral fertilizers at different doses. The BSFF increased the soil's organic matter and residual nutrient content after 42 days of the experiment, as well as the enzymatic activity of dehydrogenase and β -glucosidase, while slow mineralization rate of N afforded by BSFF resulted in the lowest yield compared to exclusive mineral fertilization and a mixture of organic/mineral fertilization [20].

On the contrary, other studies reported an increase of lettuce yield with application of several organic fertilizers from plant by-products, biochar, and olive waste derived compost. These findings highlighted the agronomic benefits of organic fertilizers if mineral N released from organic fertilizers is synchronized with the nutritional demands of lettuce crop [21,22]. Lettuce quality traits such as leaf area, and mineral, chlorophyll, and nitrate contents were

also improved with application of several organic fertilizers such as biodigested vinasse, digested livestock manure, and liquid organic fertilizer obtained from powdered soybeans, rice bran, powdered seaweed, molasses, and commercial microorganisms [23–26]. Additional studies demonstrated that soil application of organic materials like food waste compost, garlic stalk by-product, and poultry litter biochar enhanced soil enzyme activity in lettuce cultivation [22,27–31]. However, the above studies have not considered a comparison between the most promising fast-acting organic fertilizers such as poultry manure, vinasse-based fertilizer, and insect frass in greenhouse lettuce production under reduced soil fertility stemming from intensive management.

The aim of this work was: (1) to explore the dynamic of mineral N release in soil amended with three organic fertilizers such as poultry manure and vinasse-based fertilizer with the newly developed insect's frass fertilizer under controlled environment (Exp. 1—Laboratory soil incubation assay); (2) to evaluate the effects of the three tested organic fertilizers on morpho-physiological and agronomic traits of greenhouse lettuce crop and soil enzyme activity under low soil quality conditions (Exp. 2—Greenhouse trial).

2. Materials and Methods

2.1. Laboratory Soil Incubation Study (Exp. 1)

2.1.1. Experimental Conditions

An incubation test was carried out at the Horticulture Biotechnology Lab of University of Tuscia to quantify the mineral N release rate of the three organic fertilizers. Soil samples were collected at a depth of about 15 cm in the greenhouse used for the agronomic trial on lettuce for determining the N mineralization. After air-drying, greenhouse soil was screened through a 5 mm sieve. The three tested organic fertilizers in the form of dried pellets were: poultry manure (40.0 g N/kg, 17.5 g P/kg, 33.2 g K/kg; organic matter content 707 g/kg; C:N ratio 10.2), insect's frass (30.0 g N/kg, 13.1 g P/kg, 24.9 g K/kg; organic matter content 750 g/kg; C:N ratio 14.3), and a vinasse-based fertilizer containing mainly dried vinasse with the addition of poultry manure, guano and feather meal (60.0 g N/kg, 34.9 g P/kg, 124.5 g K/kg; organic matter content 500 g/kg; C:N ratio 4.8). Poultry manure (Italpollina), insect's frass (Laphrassea), and vinasse-based fertilizer (Phenix) were provided by Hello Nature Company (Rivoli Veronese, Italy). Insect's frass was derived from the digestate of black soldier fly larvae. Fertilizers were ground with a mill. Soil samples of 40 g each were prepared and mixed with each ground fertilizer at the following rate 10.0, 7.5, and 5.0 mg/g of soil for insect's frass, poultry manure, and vinasse-based fertilizer, respectively; these fertilizer rates provided the same amount of N in all tested soil:fertilizer mixtures. An unfertilized control was also included in the trial. Soil and soil:fertilizer mixtures were placed in jars (diameter 6 cm²) and watered to 70% of water container capacity.

2.1.2. Experimental Design and Analysis

The four treatments (three soil:fertilizer mixtures with insect's frass, poultry manure, and vinasse-based fertilizer, and unfertilized control) were compared in a randomized complete design with eight replicates (jars). Jars were incubated for 2 or 5 weeks in a cabinet thermostated at 24 °C in dark conditions, covering the jars with punched aluminum foil to aerate the samples. A total of 64 jars (4 treatments × 8 replicates × 2 incubation times) were included.

After 2 and 5 weeks of incubation, 8 moist samples of the soil blends for each treatment were diluted with 0.01 M CaSO₄ and stirred for 60 min before being centrifuged at 3000 × g rpm for 5 min. The supernatant was used to determine mineral N in form of nitrate (N-NO₃) [32] and ammonium (N-NH₄) [33].

Percentage of N mineralization from the amendments was calculated as the percentage of total mineral N minus the quantity from jars of nonamended soil blends divided by the total N provided by the amendment.

2.2. Greenhouse Trial (Exp. 2)

2.2.1. Experimental Conditions

The trial was carried out in a 200 m² polyethylene greenhouse at the Experimental Farm of Tuscia University (42°25' N, 12°08' E, 310 m a.s.l.) in autumn 2019. The soil was previously cultivated with vegetable crops such as melon, tomato, cucumber, and chicory. The soil contained 64% sand, 20% silt, and 16% clay. Soil chemical characteristics were: pH 7.3, electrical conductivity of saturated paste extract 5.47 dS/m, organic matter 18.8 g/kg, total nitrogen 1.16 g/kg, cation exchange capacity 25.3 meq/100g; exchangeable cations was as follow (mg/kg): K (804.1), P (15.3), Ca (3420), Mg (420), Na (502). Soil trace elements were as follow (mg/kg): soluble B (1.42), assimilable Fe (5.0), assimilable Mn (20), assimilable Cu (16.6), assimilable Zn (4.8). Based on the physico-chemical analysis, the soil was classified as sandy loam, with neutral pH, high cation exchange capacity, high salinity, and low organic matter content; moreover, the soil was very high in exchangeable K, Ca, and Mg, and available P, high in exchangeable Na, soluble B, and available Cu, medium in total N, available Mn and Zn, and low in available Fe. The above findings are in line with greenhouse soils resulting from intensive vegetable production where the intensive tillage and high temperatures lead to fast mineralization of organic matter while the lack of rainfall together with the overuse of mineral fertilizers lead to soluble salt build-up in the topsoil. The butterhead lettuce (*Lactuca sativa* L.) cultivar “Fabierto” (Rijk Zwaan, Bologna, Italy) was chosen as test crop for its early development and high productivity. The seedlings were transplanted at two-true leaf stage on 2 October in single rows (0.3 × 0.3 m) with a plant density of 11 plants m⁻². Plants were grown under natural light conditions, and the daily temperatures were maintained between 12 and 25 °C. The irrigation was performed with a drip irrigation system with in-line emitters located 0.30 m lines apart and an emitter flow rate of 3.4 L h⁻¹. Drip lines were placed in the middle of each inter-row.

2.2.2. Experimental Design, Measurements, and Analysis

The three organic fertilizers previously described [poultry manure (Italpollina), insect's frass (Laphrassea), and vinasse-based fertilizer (Phenix)] in the Exp. 1 were compared in a randomized complete block design with 4 replicates. An unfertilized control was also included. Each plot contained 4 rows of 10 plants each; the trial included a total number of 16 plots.

Organic fertilizers in the form of dried pellets were buried into the surface soil layer (0–15 cm) one day before transplant at the following rates: 2.25 t ha⁻¹ of poultry manure, 3.0 t ha⁻¹ of insect frass, and 1.5 t ha⁻¹ of vinasse-based fertiliser; these organic fertilizer rates provided the same nitrogen (N) equivalent dose of 90 kg N/ha. The organic matter (OM) and total carbon (TC) supplied by the three organic fertilizer rates were as follow (t ha⁻¹): poultry manure (1.93 OM; 0.92 TC), insect frass (2.64 OM; 1.29 TC), vinasse-based fertilizer (0.92 OM; 0.43 TC).

During the growing cycle, the Soil Plant Analysis Development (SPAD) index was measured on leaves as indicator of the chlorophyll content with the SPAD-502 instrument (Konica Minolta Europe). The readings were taken 4, 9, 15, 23, 28 days after transplanting on 20 topmost fully expanded leaves per plot. Chlorophyll fluorescence of leaves was measured with Handy-PEA (Hansatech Instruments Ltd., King's Lynn, Norfolk, UK) to determine the maximum quantum efficiency of photosystem II (Fv/Fm). The readings were taken on 20, 25, and 30 days after transplanting on 3 topmost fully expanded leaves per plot.

Lettuce plants were harvested at commercial maturity stage, after 31 days of transplanting. The yield was measured as shoot fresh weight. Shoot dry weight, root dry weight, and root to shoot ratio were determined after oven-drying plant tissues at 65 °C until constant sample weight.

Dry leaf samples harvested at the end of the trials were ground separately in a Wiley mill to pass through a 20-mesh screen, then 0.5 g of the dried plant tissues were analyzed for the following mineral elements: N, P, K, Ca, Mg, Fe, Mn, Zn, Cu, Ni, Na, Al, Ba, Cd, Co, Cr, Pb, Se, and Sn. Nitrogen concentration in the plant tissues was determined after mineralization with sulfuric acid by 'Kjeldahl method' [34] while the other elements were

determined by dry ashing at 400 °C for 24 h, dissolving the ash in 1:20 HNO₃, and assaying the solution obtained using an inductively coupled plasma emission spectrophotometer (ICP Iris; Thermo Optek, Milano, Italy) [35].

At the end of the experiment, soil samples were randomly collected in the topsoil (0–0.15 m) within each plot by using a metal soil probe (2.5 cm diameter). Ten cores were collected from each plot and combined into one composite sample and homogenized through a 2 mm sieve. Soil samples were refrigerated at 4 °C and transported to laboratory to determine dry weight, active carbon, and soil enzyme activity. Each analysis was carried out on two laboratory replicates.

Five grams of fresh soil was weighed, and oven dried at 105 °C until constant weight to determine soil dry weight. Soil samples were air-dried and further sieved (1 mm) before labile soil carbon determination [36]. According to the method, the KMnO₄ stock solution (0.2 M) was diluted 1:10 with distilled water and mixed to soil samples (20 mL of solution and 5 g of dried soil). The soil KMnO₄ mixture was agitated for 2 min at 120 rpm and then allowed to stand for 10 min. Once the soil was well deposited, 0.50 mL of supernatant was added to 45 mL distilled water and shaken 10 s. Each diluted solution was then poured into an optically calibrated glass vial and the light absorption was measured at a 550 nm fixed wavelength using a UV-Vis Shimadzu, UV-1800 spectrophotometer (Shimadzu Corporation, Kyoto, Japan).

Acid phosphatase (AcP, EC 3.1.3.2) and, alkaline phosphatase (AlkP, EC 3.1.3.1) activities were assayed by the hydrolysis rate of *p*-nitrophenyl disodium orthophosphate (*p*-NPP) supplied as substrate [37,38]. Arylsulphatase (ArS, EC 3.1.6.1) was assayed using *p*-nitrophenyl sulphate (*p*-NPS) as substrate. Dehydrogenase (DHA, EC 1.1.1) activity was assayed using tetrazolium salt as substrate [39] and N-acetyl-β-D-glucosaminidase (NAGase, EC 3.2.1.30; EC 3.2.1.52) activity was determined using *p*-nitrophenyl-N-acetyl-β-D-glucosaminidine as substrate [40]. In addition, total hydrolytic activity (THA) was assessed by the fluorescein diacetate method that measures activities of several enzymes such as proteases, lipases and non-specific esterases, related to organic matter decomposition [41]. All spectrophotometric measurements were carried out by UV-1800 Shimadzu spectrophotometer. The enzyme activities were expressed as µg of product per gram of oven-dry weight soil per hour.

2.3. Statistical Analysis

All data were subjected to ANOVA using the SPSS22 software package (Chicago, IL, USA). Means were separated using Tukey's range test performed at a 5% level of significance. For each enzyme, Pearson correlation was performed between enzyme activity and organic fertilizer rate. A hierarchical cluster analysis on the morpho-physiological crop traits, soil enzymatic activity and soil active carbon was performed, and a heatmap was generated using the ClustVis online tool. Matrix values were normalized as $\ln(x + 1)$, with Euclidean distance and complete linkage.

3. Results

3.1. Nitrogen Mineralized from Organic Fertilizers during Laboratory Incubation Assay

To determine the nitrogen release pattern from poultry manure, insect frass and vinasse-based fertilizers, a laboratory aerobic incubation experiment was conducted for 5 weeks, and the outcomes are shown in Figure 1. After 2 weeks of incubation, vinasse-based fertilizer had highest N mineralization followed by poultry manure and insect frass. After 5 weeks of incubation, only vinasse-based fertilizer had higher significant N mineralization compared to insect frass while poultry manure has an intermediate value.




Figure 1. Mean cumulative N mineralization by fertilizer type at 2 and 5 weeks of aerobic incubation of soil: fertilizer blends. Mean square error at 2 wk = 0.53; Mean square error at 5 wk = 145.85; Coefficient of variation at 2 wk = 46.8%; Coefficient of variation at 5 wk = 26.9%; Tukey's least significant difference (5%) at 2 wk = 0.92; Tukey's least significant difference (5%) at 5 wk = 15.22. Different letters within each week of incubation indicate significant differences according to Tukey's range test (5%).

3.2. Agronomic Response of Lettuce and Soil Enzymatic Activity under Greenhouse Trial

Organic fertilizers significantly ($p < 0.001$) affected chlorophyll concentration, expressed as SPAD index, of lettuce leaves at 9 and 15 days after transplanting (DAT) while no significant differences were found among treatments at 4 DAT (Table 1). At 9 and 15 DAT, the highest SPAD index was recorded in plant leaves treated with vinasse-based fertilizer followed by insect frass and poultry manure which showed similar values; the lowest SPAD value was recorded in untreated control.

Table 1. Effect of fertilizer treatments on chlorophyll concentration expressed as SPAD index of lettuce leaves.

Fertilizer Treatment	SPAD Index				
	4 DAT	9 DAT	15 DAT	23 DAT	28 DAT
Control	14.93	19.52 c	24.67 c	24.90	24.65
Poultry manure	14.53	23.30 b	26.27 b	24.31	24.67
Insect frass	15.46	23.13 b	26.33 b	24.38	24.88
Vinasse-based fertilizer	15.25	24.63 a	28.29 a	25.20	25.24
MSE	0.35	0.39	0.41	0.64	1.13
CV (%)	4.27	8.99	5.47	3.28	3.26
Tukey (5%)	1.25	1.32	1.34	1.68	2.24

MSE = mean square error; CV = coefficient of variation; Tukey (5%) = Tukey's least significant difference (5%). Different letters within each column indicate significant differences according to Tukey's range test (5%).

The outcomes of the effect of organic fertilizers on maximum quantum yield of PSII (F_v/F_m) of lettuce leaves revealed that the application of the three organic fertilizers had no effect on F_v/F_m at 20 and 30 DAT while a significant ($p < 0.05$) decrease of F_v/F_m value was recorded at 25 DAT with poultry manure application in comparison to other organic treatments and unfertilized control (Table 2).

Table 2. Effect of fertilizer treatments on maximum quantum yield of PSII (Fv/Fm) of lettuce leaves.

Fertilizer Treatment	Fv/Fm		
	20 DAT	25 DAT	30 DAT
Control	0.837	0.831 a	0.852
Poultry manure	0.836	0.822 b	0.845
Insect frass	0.842	0.834 a	0.855
Vinasse-based fertilizer	0.838	0.830 ab	0.849
MSE	0.000279	0.000014	0.000465
CV (%)	1.804	0.577	2.313
Tukey (5%)	0.035	0.008	0.045

MSE = mean square error; CV = coefficient of variation; Tukey (5%) = Tukey's least significant difference (5%). Different letters within each column indicate significant differences according to Tukey's range test (5%).

The results of the effect of fertilizer treatments on morpho-physiological and agronomic traits of lettuce crop are shown in Table 3. The application of the three organic fertilizers affected significantly both the shoot fresh ($p < 0.001$) and dry weight ($p < 0.01$) while no significant differences were found on the root dry weight and the root to shoot ratio (Table 3). The highest value of shoot fresh weight was obtained in plants treated with vinasse-based fertilizer and insect frass in comparison with the control while the poultry manure treatment gave intermediate values.

Table 3. Effect of fertilizer treatment on shoot fresh weight, shoot and root dry weight, and root to shoot ratio of lettuce plants.

Fertilizer Treatment	Shoot Fresh Weight (g Plant ⁻¹)	Shoot Dry Weight (g Plant ⁻¹)	Root Dry Weight (g Plant ⁻¹)	Root to Shoot Ratio
Control	132.31 c	5.13 b	1.91	0.39
Poultry manure	172.99 b	7.49 ab	2.46	0.34
Insect frass	230.99 a	9.97 a	2.41	0.24
Vinasse-based fertilizer	233.15 a	9.33 a	2.70	0.29
MSE	331.5	2.01	0.22	0.01
CV (%)	24.21	29.05	21.80	31.42
Tukey (5%)	38.24	2.98	1.00	0.19

MSE = mean square error; CV = coefficient of variation; Tukey (5%) = Tukey's least significant difference (5%). Different letters within each column indicate significant differences according to Tukey's range test (5%).

Among the tested organic fertilizers, insect's frass significantly ($p < 0.05$) reduced the nitrate concentration in lettuce leaves in comparison with the other organic fertilizers while untreated control gave intermediate values which were not significantly different from the organic fertilizer treatments (Figure 2).

Except for the leaf P concentration which was significantly ($p < 0.01$) highest in vinasse-based fertilizer treated plants compared to the other fertilizer treatments and control, the soil amendment with the three organic fertilizers did not affect the macronutrient concentrations in lettuce leaves (Table 4).

Apart from for Mo, which was significantly ($p < 0.01$) lowest in vinasse-based treatment, no significant differences were recorded for the other micronutrients in lettuce leaves (Table 5).




Figure 2. Effect of fertilizer treatments on nitrate concentration in lettuce leaves. Mean square error = 292,554; Coefficient of variation = 20.5%; Tukey's least significant difference (5%) = 1135.9. Different letters indicate significant differences according to Tukey's range test (5%).

Table 4. Effect of fertilizer treatments on macronutrient concentration in lettuce leaves.

Fertilizer Treatment	Macronutrients (g/kg DW)				
	N	P	K	Ca	Mg
Control	44.16	3.33 b	82.72	14.09	5.63
Poultry manure	43.10	4.05 b	90.30	12.93	5.95
Insect frass	41.05	4.07 b	84.36	12.51	5.78
Vinasse-based fertilizer	42.55	5.09 a	85.52	11.80	5.51
MSE	3.09	0.20	40.67	1.78	1.29
CV (%)	4.57	18.51	7.47	11.45	18.02
Tukey (5%)	3.69	0.95	13.39	2.80	2.39

MSE = mean square error; CV = coefficient of variation; Tukey (5%) = Tukey's least significant difference (5%). Different letters within each column indicate significant differences according to Tukey's range test (5%).

Table 5. Effect of fertilizer treatments on essential trace elements in lettuce leaves.

Fertilizer Treatment	Essential Trace Elements (mg/kg DW)						
	B	Fe	Mn	Zn	Cu	Mo	Ni
Control	21.72	190.89	285.36	52.40	12.78	0.37 a	0.29
Poultry manure	26.70	185.91	314.19	65.55	19.77	0.39 a	0.20
Insect frass	25.18	196.72	297.24	55.85	17.98	0.36 a	0.27
Vinasse-based fertilizer	25.14	128.06	254.83	51.59	14.52	0.25 b	0.18
MSE	16.44	9511.57	4866.77	177.74	17.32	0.002	0.011
CV (%)	16.55	52.32	23.02	23.48	28.81	19.31	44.78
Tukey (5%)	8.51	204.81	146.50	28.00	8.74	0.08	0.22

MSE = mean square error; CV = coefficient of variation; Tukey (5%) = Tukey's least significant difference (5%). Different letters within each column indicate significant differences according to Tukey's range test (5%).

The effect of fertilizer treatments on non-essential trace elements in lettuce leaves was significant for the heavy metals Co and Pb while there were no significant differences for Na, Al, Ba, Cd, Cr, Se and Sn (Table 6). The vinasse-based fertilizer was able to reduce significantly ($p < 0.01$) the amount of Co in lettuce leaves compared to the control which had the highest value. Moreover, the soil amendment with the three organic fertilizers leads to a significant ($p < 0.01$) reduction in leaf Pb concentration in comparison to unfertilized control.

Table 6. Effect of fertilizer treatments on non-essential trace elements in lettuce leaves.

Fertilizer Treatments	Non-Essential Trace Elements (mg/kg DW)								
	Na	Al	Ba	Cd	Co	Cr	Pb	Se	Sn
Control	3.62	440.17	19.04	0.24	0.23 a	0.28	1.29 a	0.92	0.89
Poultry manure	3.83	402.93	22.28	0.36	0.17 ab	0.32	0.81 b	0.84	0.83
Insect frass	4.66	338.47	25.35	0.26	0.17 ab	0.40	0.77 b	0.98	0.83
Vinasse-based fertilizer	3.61	254.33	17.63	0.28	0.09 b	0.25	0.52 b	1.02	0.83
MSE	1.127	32,475.66	42.56	0.004	0.001	0.027	0.044	0.034	0.004
CV (%)	26.68	49.27	31.32	25.99	37.12	50.66	40.59	19.04	7.48
Tukey (5%)	2.23	378.44	13.70	0.13	0.08	0.35	0.44	0.39	0.14

MSE = mean square error; CV = coefficient of variation; Tukey (5%) = Tukey's least significant difference (5%). Different letters within each column indicate significant differences according to Tukey's range test (5%).

Soil active carbon at the end of the trial was significantly ($p < 0.001$) affected by fertilizer treatments ($p < 0.001$) with higher values with poultry manure (444.3 mg C/kg) and insect's frass (485.8 mg C/kg) in comparison with vinasse-based fertilizer (375.7 mg C/kg) and unfertilized control (369.5 mg C/kg).

In the current experiment, the application of poultry manure and insect's frass significantly increased the AcP ($p < 0.001$), AlkP ($p < 0.001$) and DHA ($p < 0.001$) in comparison to the control and vinasse-based fertilizer, which were not significantly different (Table 7). The ArS was significantly ($p < 0.01$) highest after the application of insect's frass, followed by poultry manure and vinasse-based fertilizer which had similar values while the lowest value was observed in the unfertilized control. The NAGase—Chitinase had a significantly ($p < 0.001$) higher value in insect's frass treatment in comparison to the control and vinasse-based fertilizer treatment while no significant difference was recorded with poultry manure application. In case of applying vinasse-based fertilizer, NAGase—Chitinase had significantly higher value than unfertilized control. THA had significant ($p < 0.001$) higher values in insect's frass and poultry manure than in control while vinasse-based fertilizer had an intermediate value (Table 7). A significant Pearson correlation was observed between organic fertilizer rate and soil enzyme activity for AcP ($r = 0.981^*$), ArS ($r = 0.938^*$), DHA ($r = 0.933^*$), and THA ($r = 0.955^*$). Moreover, no significant correlation between organic fertilizer rate and soil enzyme activity was observed for AlkP and NAGase—chitinase.

Hierarchical clustering analysis coupled with a heatmap was performed to provide a visual representation of the changes in morpho-physiological crop traits, soil active carbon, and soil enzymatic activity after the application of the three organic fertilizer treatments (Figure 3). In the treatment dendrogram the unfertilized control was clearly separated from the other fertilizer treatments due to the decrease of shoot and root growth, leaf SPAD index at 9 and 15 days after transplanting (DAT), leaf concentration of P, K, Mg, B, Cu, Cd, soil enzymatic activity (AcP, AlkP, ArS, DHA, NAGase, THA), and soil active C, and the increase of leaf concentration of N, Pb, Ca, Co, Sn and root to shoot ratio. Moreover, vinasse-based fertilizer treatment was clearly separated from the two other organic fertilizers due to the increase of root dry weight, leaf SPAD index at 9, 15, 23, and 28 DAT and leaf concentration of P, and the decrease of leaf concentration of Ca, Mg, Fe, Mn, Zn, Cu, Mo, Al, Ba, Cr, Na, Pb, Co, Ni, soil enzymatic activity (AcP, AlkP, ArS, DHA, NAGase, THA) and soil active C. Insect frass and poultry manure were clustered together on the same branch; the main differences between these two organic fertilizers were due to the increase of shoot fresh and dry weight, maximum quantum yield of PSII (Fv/Fm) of leaves at 20, 25, and 30 DAT,

leaf SPAD index at 4 DAT, and leaf concentration of Se, Cr, Ba, Na, Ni, and the decrease of root to shoot ratio, leaf concentration of N, NO₃, K, Zn, and Cd in insect frass compared to poultry manure treatment.

Table 7. Effect of fertilizer treatment on soil enzyme activity per gram of dry soil per hour in greenhouse lettuce trial.

Fertilizer Treatment	AcP ($\mu\text{g } p\text{-Nitrophenol/g h}$)	AlkP ($\mu\text{g } p\text{-Nitrophenol/g h}$)	ArS ($\mu\text{g } p\text{-Nitrophenol/g h}$)	DHA ($\mu\text{g Triphenyl formazan/g h}$)	NAGase— Chitinase ($\mu\text{g } p\text{-Nitrophenol/g h}$)	THA ($\mu\text{g Fluorescein/g h}$)
Control	48.74 b	362.25 b	4.83 c	0.42 b	8.84 c	14.26 c
Poultry manure	91.61 a	619.94 a	8.29 b	1.87 a	47.81 ab	63.93 a
Insect frass	99.35 a	561.58 a	11.43 a	1.88 a	57.61 a	61.31 a
Vinasse-based fertilizer	65.35 ab	336.50 b	8.21 b	0.84 b	30.85 b	35.56 b
MSE	298.53	1766.50	1.81	0.19	71.26	26.31
CV (%)	34.13	28.14	32.90	61.52	56.66	49.10
Tukey (5%)	36.28	88.26	2.83	0.93	17.73	10.77

AcP = acid phosphatase; AlkP = alkaline phosphatase, ArS=arylsulfatase, N-acetyl- β -D-glucosaminidase (NAGase-chitinase); DHA = dehydrogenase activity; THA = total hydrolase activity. MSE = mean square error; CV = coefficient of variation; Tukey (5%) = Tukey's least significant difference (5%). Different letters within each column indicate significant differences according to Tukey's range test (5%).




Figure 3. Cluster heat map analysis summarizing lettuce plants response to organic fertilizer application. Original values are $\ln(x + 1)$ -transformed. Columns and rows are clustered using Euclidean distance and complete linkage.

4. Discussion

The rate and timing of mineralization of organic fertilizer is a key factor to meet the plant's needs for nutrients. Mineralization process of organic fertilizers is affected by several factors, such as quality of organic matter, soil temperature, soil water availability, etc. Among these factors, C/N ratio of organic fertilizer plays a pivotal role in mineralization and immobilization processes of N in soils. The lower the C/N ratio, the faster nitrogen is supplied into the soil for direct crop absorption [42]. Our data showed in line with several authors [43–45], that vinasse-based fertilizer had the highest N mineralization rate. These results can be linked to its lower C/N ratio (4.8) in comparison to poultry manure (10.2) and insect frass (14.3). Therefore, vinasse-based fertilizer is a more suitable fertilizer than poultry manure and insect' frass in terms of supplying the mineral N in soil during the short crop cycle of lettuce under greenhouse conditions.

Leaf chlorophyll concentration has been linked to plant stress and nutrient status. As a component of chlorophyll molecule, nitrogen is a major nutrient affecting leaf chlorophyll concentration. The SPAD meter was demonstrated to be a useful tool for non-destructively assessing leaf chlorophyll concentration. In the greenhouse trial, leaf SPAD index at 4 DAT was not significantly affected by treatments, indicating that a significant proportion of absorbed nitrogen during the early stages of plant growth may have been used to generate structural elements of the plant rather than chlorophyll as suggested by several authors [46]. Moreover, the significant difference in chlorophyll concentration at 9 and 15 DAT could be related to the faster N mineralization of vinasse-based fertilizer in comparison to other treatments and to the high plant demand of nitrogen during the exponential phase of plant growth [47]. On the other hand, the non-significance among all treatments at 23 and 28 DAT could be explained by the low N demand by the lettuce plants during this final growth stage.

Maximum quantum yield of PSII (F_v/F_m) is a sensitive indicator of plant photosynthetic ability of leaves where low F_v/F_m values may indicate downregulation of photosynthesis resulting from stress and/or photoinhibition [48,49]. In this current study, the F_v/F_m values observed during the growing cycle in all treatments were in the range of 0.75 to 0.85 reported for healthy leaf crops [50,51].

According to the FAO data of crop salt tolerance, lettuce is considered a moderately sensitive crop to salinity with a salinity threshold value for shoot fresh weight of 1.3 dS/m and a percentage of shoot yield reduction per dS/m above the threshold value of 13. Considering that the soil EC of saturated paste extract was 5.47 dS/m, the relative crop yield reduction determined using the Maas–Hoffman model [52] was 45.8% indicating a strong negative impact of soil salinity on shoot fresh biomass in the experimental plots. Soil amendment with organic fertilizers increased shoot fresh weight of lettuce especially in vinasse-based and insect's frass treatments. Shoot dry weight showed a similar behavior of shoot fresh weight, indicating that the increase of shoot fresh weight under organic fertilization regimes in comparison with unfertilized control was due to an enhancement of carbon assimilation. This finding indicates an important role of the three organic fertilizers in mitigating the negative impact of salinity on lettuce growth in comparison to the unfertilized control.

Organic matter can be incorporated into soil to provide a positive impact on the physical, chemical, and biological characteristics of the soil [53–56]. These benefits are mainly ascribed to the supply of organic compounds and especially humic substances through the organic amendment. Several authors reported that humic substances can increase aggregate stability in soil [57], stimulate secondary metabolism pathways involved in stress response [58], immobilize heavy metals by binding or redox reaction [59–61], stimulate the plant growth, biosynthetic activity, and nutrient uptake [62,63] activate antioxidative enzymatic function, and control the reactive oxygen species content in plants [64]. Moreover, organic fertilization can promote beneficial microflora as the result of the microbial cells' ability to adapt to osmotic stress by creating osmolytes, which counteract osmotic stress, thanks to the availability of carbon from the additional organic matter [53,65,66].

Nitrate concentration of lettuce leaves is an important quality trait affecting marketability of product. The Commission Regulation EU No 1258/2011 [67] amending Regulation EC No 1881/2006 established maximum levels for nitrates in lettuce grown in greenhouse from 1st October until 31st March to be 5000 mg kg⁻¹ of fresh weight. In the current study, nitrate concentrations in lettuce leaves were always lower than the EU threshold value indicating no restriction for commercialization. However, insect's frass treatment gave a lower leaf nitrate concentration in comparison with the other treatments, indicating a positive role of insect's frass in the activation of key enzymes for nitrogen assimilation in plant cells (e.g., nitrate reductase).

Organic matter is the major component of organic fertilizer which increases the soil's macronutrient and micronutrient concentrations, and through enhancing its structure, it leads to increased soil fertility and quality [68]. Considering the effect of organic fertilizers on macro and micronutrients concentration in lettuce leaves, significant differences were recorded only on leaf concentration of P and Mo, which were the highest and the lowest in vinasse-based treatment, respectively. The increase of leaf P concentration in vinasse-based treatment can be related to the different P supplied to the plants which was higher with vinasse-based fertilizer (52.4 kg P ha⁻¹) than poultry manure and insect's frass (both 39.3 kg P ha⁻¹) and untreated control. Moreover, the fast mineralization of vinasse-based fertilizer may also have enhanced the soil P availability for plant uptake. According to the sufficiency range for butterhead lettuce under greenhouse conditions [69], the macronutrients and micronutrients concentrations in lettuce leaves were in the sufficient ranges as for N, Mg, Fe, Zn, and Mo, while they exceeded the sufficient range as for K, Ca, Mn, and Cu. Leaf P concentrations in all treatments were below the sufficient range reported by the authors [69], indicating the P availability was probably a critical factor in plant growth under the current experiment.

Regarding the effect of organic fertilizers on non-essential trace elements in lettuce leaves, the reduction of leaf Co concentration by vinasse-based fertilizer application and the decrease of leaf Pb concentration by the soil amendment with the three organic fertilizers indicate a positive role of organic matter in mitigating the accumulation of some heavy metals in plant tissues due to adsorption and chelating actions of heavy metals ions [70–72]. Heavy metals are immobilized by humic substances primarily by binding (complexation, ion exchange, and physical adsorption) or redox reaction [59–61]. Binding heavy metals can fix and limit their fluidity in the soil, whereas heavy metal redox reactions can diminish or remove their toxicity [73]. Moreover, plant uptake of heavy metals can be restricted by the increasing of microorganisms in organically fertilized soil [70,71,74,75]. Similar results were reported in an incubation experiment where amendment with organic fertilizers was effective in reducing the availability of polluted soil with the heavy metal Cd, and the inhibitory effect was stronger as the quantity of organic fertilizer was augmented [71].

Soil enzymes are mostly microbial in origin and are tightly related to microbial abundance and activity. They are critical in catalyzing reactions required for organic matter decomposition and nutrient cycling [76]. Climate conditions, type of amendment, cultivation methods, crop type, and edaphic qualities are all factors that influence enzyme catalyzed reactions. Soil enzyme activities are mainly influenced by organic matter content [77] and are frequently used as indicators of microbial activity and soil fertility [78]. In this study, soil enzymes responsible for the major nutrient cycles (N, P, C and S) were chosen. Phosphatases (AcP, AlkP) are very useful enzymes in agriculture because they hydrolyze organic phosphorus compounds and convert them into inorganic phosphate that plants can absorb [79]. Arylsulfatase (ArS) is an enzyme that participates in the hydrolysis of arylsulphate esters by the fission of the oxygen-sulfur (O-S) bond. This enzyme is known to play a role in the mineralization of ester sulfate in soils [80]. Furthermore, because only fungi (not bacteria) possess ester sulfate, the substrate of arylsulfatase [81,82], it may be an indirect indication of fungal abundance in the soil. Dehydrogenases (DHA) are among the most significant soil enzymes for determining biological activity in the soil because they represent the entire oxidative metabolic activity of soil microbial community [83]. N-acetyl-β-D-glucosaminidases (NAGase—Chitinase) are abundant in nature, and the low molecular

weight amino sugars produced by their hydrolysis constitute an essential source of nitrogen for soil microbes. β -glucosidase catalyzes the hydrolysis of β -D-glucopyranosides and is one of three or more enzymes necessary for the saccharification of cellulose [84,85]. The total hydrolytic activity (THA) measurement is a simple method for assessing total microbial activity [41]. Several authors have reported an increase in soil microbial biomass, bacteria populations, fungi and actinomycetes, as well as soil enzyme activities after applying organic fertilizers such as manures and composts to plants [6,8,27,28,86]. In agreement with the above findings, soil enzyme activities increased in organically treated plots especially with the application of poultry manure and insect's frass. The soil enzymatic activity for AcP, ArS, DHA, and THA was positively correlated with the organic fertilizer rate while no significant correlation between organic fertilizer rate and soil enzymatic activity was observed for AlkP and NAG-ase—chitinase, indicating that the quality of organic matter supplied by fertilizers may have differentially affected their activity. Several authors found that the quality of organic matter may alter the organization of the soil microbial population because microorganisms have substrate preferences [87]; the different structure of microbial community resulting from organic fertilizer application can explain the variation of the soil enzyme activity and the lack of significant correlation for AlkP and NAG-ase.

The available or labile portion of soil organic matter, also known as active carbon, is a sensitive tool for measuring soil carbon change. Active carbon is used as a readily available food and energy source for the soil microorganisms. At the end of the trial, organic fertilization with poultry manure or insect's frass was able to raise the soil active carbon in comparison to vinasse-based fertilizer and unfertilized control. It has been demonstrated that soil amendment with high C/N ratio may improve and maintain soil labile C for a longer period [88]. Therefore, the highest C/N ratio of poultry manure and insect's frass together with the highest fertilizer rate can explain the increase of soil active carbon at the end of the trial compared to vinasse-based fertilizer.

5. Conclusions

Sustainable intensification of greenhouse vegetable production relies on the application of organic matter to maintain soil fertility and support plant nutrition providing that nutrient release in available forms for plant uptake match the crop nutrient requirements. Based on the laboratory incubation assay, the vinasse-based fertilizer can be recommended for supporting plant nutrition due to the fast mineralization and release of mineral N which is highly required during the first growing period of the short lettuce cultivation cycle. Our data demonstrated that application of organic fertilizers such as poultry manure, insect's frass, and vinasse-based fertilizer in greenhouse lettuce production was a good strategy for reducing stressful conditions arising from the soil salinization and soil organic matter depletion due to intensive vegetable crop production. Organic fertilizers enhanced lettuce growth as demonstrated by the increase of shoot fresh and dry weight. Moreover, insect frass was also effective in improving the lettuce quality by reducing the nitrate concentration in leaves. In terms of soil quality, poultry manure and insect's frass were more effective in restoring soil biological fertility as demonstrated by the increase of soil active carbon and the enzyme activity involved in nutrient cycling.

Finally, further research is needed to investigate the potential benefits of combined applications of vinasse-based fertilizer and the other organic fertilizers for supporting plant nutrition and at the same time enhancing the beneficial effects on soil fertility in lettuce production.

Author Contributions: Conceptualization, M.C. and G.C.; methodology, M.C., P.I., P.B. and G.C.; validation, M.C., A.E.C., P.I., P.B. and G.C.; formal analysis, M.C., P.I., Y.R. and G.C.; investigation, M.C., P.I., P.B. and G.C.; resources, M.C., P.I. and G.C.; data curation, M.C., A.E.C., P.I., Y.R. and G.C.; writing—original draft preparation, M.C., A.E.C. and G.C.; writing—review and editing, M.C., A.E.C., Y.R., P.B. and G.C.; visualization, M.C., A.E.C. and G.C.; supervision, M.C. and G.C.; project administration, M.C. and G.C.; funding acquisition, G.C. All authors have read and agreed to the published version of the manuscript.

Funding: This work was partially funded by MIUR in the frame of the initiative “Departments of excellence”, Law 232/2016.

Data Availability Statement: Data available on request from the corresponding author.

Acknowledgments: We thank Antonio Fiorillo and Rares Balan for their assistance in supporting greenhouse and laboratory assays. We also thank Hello Nature for providing the tested organic fertilizers.

Conflicts of Interest: The authors declare no conflict of interest.

References

- Hamidov, A.; Helming, K.; Bellocchi, G.; Bojar, W.; Dalgaard, T.; Ghaley, B.B.; Hoffmann, C.; Holman, I.; Holzkämper, A.; Krzeminska, D.; et al. Impacts of climate change adaptation options on soil functions: A review of European case-studies. *Land Degrad. Dev.* **2018**, *29*, 2378–2389. [CrossRef] [PubMed]
- Sonneveld, C. Effects of Salinity on Substrate Grown Vegetables and Ornamentals in Greenhouse Horticulture. Ph.D. Thesis, University of Wageningen, Wageningen, The Netherlands, 2000.
- European Commission. *Farm to Fork: New Rules to Reduce the Risk and Use of Pesticides in the EU*; European Commission: Brussels, Belgium, 2022; pp. 1–3. Available online: https://ec.europa.eu/commission/presscorner/detail/en/qanda_22_3694 (accessed on 14 October 2022).
- Willer, H.; Trávní, J.; Meier, C.; Schlatter, B. (Eds.) *The World of Organic Agriculture Statistics and Emerging Trends 2022*; Research Institute of Organic Agriculture FiBL, Frick, and IFOAM—Organics International: Bonn, Germany, 2022; pp. 1–345. Available online: <https://www.fibl.org/fileadmin/documents/shop/1344-organic-world-2022.pdf> (accessed on 14 October 2022).
- Sun, X.; Ye, Y.; Ma, Q.; Guan, Q.; Jones, D.L. Variation in enzyme activities involved in carbon and nitrogen cycling in rhizosphere and bulk soil after organic mulching. *Rhizosphere* **2021**, *19*, 100376. [CrossRef]
- Antonious, G.F.; Turley, E.T.; Dawood, M.H. Monitoring Soil Enzymes Activity before and after Animal Manure Application. *Agriculture* **2020**, *10*, 166. [CrossRef]
- Hu, W.; Jiao, Z.; Wu, F.; Liu, Y.; Dong, M.; Ma, X.; Fan, T.; An, L.; Feng, H. Long-term effects of fertilizer on soil enzymatic activity of wheat field soil in Loess Plateau, China. *Ecotoxicology* **2014**, *23*, 2069–2080. [CrossRef] [PubMed]
- Chang, E.-H.; Chung, R.-S.; Wang, F.-N. Effect of different types of organic fertilizers on the chemical properties and enzymatic activities of an Oxisol under intensive cultivation of vegetables for 4 years. *Soil Sci. Plant Nutr.* **2008**, *54*, 587–599. [CrossRef]
- Ciriminna, R.; Scurria, A.; Tizza, G.; Pagliaro, M. Volcanic ash as multi-nutrient mineral fertilizer: Science and early applications. *JSA Rep.* **2022**, *2*, 528–534. [CrossRef]
- Yang, L.; Zhang, Y.; Li, F. Soil Enzyme Activities and Soil Fertility Dynamics. In *Advances in Citrus Nutrition*; Springer: Dordrecht, The Netherlands, 2012; pp. 143–156. [CrossRef]
- Tittarelli, F.; Bâth, B.; Ceglie, F.; García, M.; Möller, K.; Reents, H.; Védie, H.; Voogt, W. Soil fertility management in organic greenhouse: An analysis of the European context. *Acta Hortic.* **2017**, *1164*, 113–126. [CrossRef]
- Loecke, T.D.; Cambardella, C.A.; Liebman, M. Synchrony of net nitrogen mineralization and maize nitrogen uptake following applications of composted and fresh swine manure in the Midwest U.S. *Nutr. Cycl. Agroecosyst.* **2012**, *93*, 65–74. [CrossRef]
- Cardarelli, M.; Salerno, A.; Rea, E.; Colla, G. Nitrogen availability from organic and organic-mineral fertilizers. In Proceedings of the 18th International Symposium of CIEC ‘More Sustainability in Agriculture: New Fertilizers and Fertilization Management’, Rome, Italy, 8–12 November 2009.
- Carassay, L.R.; Bustos, D.A.; Golberg, A.D.; Taleisnik, E. Tipburn in salt-affected lettuce (*Lactuca sativa* L.) plants results from local oxidative stress. *J. Plant Physiol.* **2012**, *169*, 285–293. [CrossRef]
- Sago, Y. Effects of Light Intensity and Growth Rate on Tipburn Development and Leaf Calcium Concentration in Butterhead Lettuce. *HortScience* **2016**, *51*, 1087–1091. [CrossRef]
- Pavlou, G.C.; Ehalotis, C.D.; Kavvadias, V.A. Effect of organic and inorganic fertilizers applied during successive crop seasons on growth and nitrate accumulation in lettuce. *Sci. Hortic.* **2007**, *111*, 319–325. [CrossRef]
- Gunes, A.; Post, W.N.K.; Kirkby, E.A.; Aktas, M. Influence of partial replacement of nitrate by amino acid nitrogen or urea in the nutrient medium on nitrate accumulation in NFT grown winter lettuce. *J. Plant Nutr.* **1994**, *17*, 1929–1938. [CrossRef]
- Colla, G.; Kim, H.-J.; Kyriacou, M.C.; Rouphael, Y. Nitrate in fruits and vegetables. *Sci. Hortic.* **2018**, *237*, 221–238. [CrossRef]
- Craddock, V.M. Toxicology: Nitrosamines and human cancer: Proof of an association? *Nature* **1983**, *306*, 638. [CrossRef]
- Esteves, C.; Fareleira, P.; Castelo-Branco, M.; Lopes, I.; Mota, M.; Murta, D.; Menino, R. Black soldier fly larvae frass increases the soil’s residual nutrient content and enzymatic activity—A lettuce production trial. *J. Insects Food Feed* **2022**, *8*, 1431–1440. [CrossRef]
- Cavalheiro, T.R.T.; Alcoforado, R.D.O.; Silva, V.S.D.A.; Coimbra, P.P.S.; Mendes, N.D.S.; Cavalcanti, E.D.C.; Jurelevicius, D.D.A.; Gonçalves, C.B.D.A. The Impact of Organic Fertilizer Produced with Vegetable Residues in Lettuce (*Lactuca sativa* L.) Cultivation and Antioxidant Activity. *Sustainability* **2020**, *13*, 128. [CrossRef]
- Trupiano, D.; Coccozza, C.; Baronti, S.; Amendola, C.; Vaccari, F.P.; Lustrato, G.; Di Leonardo, S.; Fantasma, F.; Tognetti, R.; Scippa, G.S. The Effects of Biochar and Its Combination with Compost on Lettuce (*Lactuca sativa* L.) Growth, Soil Properties, and Soil Microbial Activity and Abundance. *Int. J. Agron.* **2017**, *2017*, 3158207. [CrossRef]

23. Santos, C.M.; Gonçalves, E.R.; Endres, L.; Gomes, T.C.A.; Jadoski, C.J.; Nascimento, L.A.; Santos, E.D. Photo-synthetic measurements in lettuce submitted to different agroindustrial residue composting. *Pesqui. Apl. Agrotecnol.* **2010**, *3*, 103–112. [\[CrossRef\]](#)
24. Ueno, C.R.J.; Da Costa, A.C.S.; Gimenes, M.L.; Zanin, G.M. Agricultural recycling of biodigested vinasse for lettuce production. *Ambient. E Agua Interdiscip. J. Appl. Sci.* **2014**, *9*, 593–601. [\[CrossRef\]](#)
25. Trinchera, A.; Baratella, V.; Rinaldi, S.; Renzaglia, M.; Marcucci, A.; Rea, E. Greenhouse lettuce: Assessing nutrient use efficiency of digested livestock manure as organic N-fertiliser. *Acta Hort.* **2014**, *1041*, 63–69. [\[CrossRef\]](#)
26. Liu, C.-W.; Sung, Y.; Chen, B.-C.; Lai, H.-Y. Effects of Nitrogen Fertilizers on the Growth and Nitrate Content of Lettuce (*Lactuca sativa* L.). *Int. J. Environ. Res. Public Health* **2014**, *11*, 4427–4440. [\[CrossRef\]](#) [\[PubMed\]](#)
27. Lee, J. Effect of food waste compost on microbial population, soil enzyme activity and lettuce growth. *Bioresour. Technol.* **2004**, *93*, 21–28. [\[CrossRef\]](#)
28. Akça, M.O.; Namlı, A. Effects of poultry litter biochar on soil enzyme activities and tomato, pepper and lettuce plants growth. *Eurasian J. Soil Sci.* **2015**, *4*, 161–168. [\[CrossRef\]](#)
29. Han, X.; Cheng, Z.; Meng, H. Growth of wheat and lettuce and enzyme activities of soils under garlic stalk decomposition for different durations: Function and utilization of garlic stalk decomposition for different durations. *J. Sci. Food Agric.* **2017**, *97*, 2727–2735. [\[CrossRef\]](#) [\[PubMed\]](#)
30. Han, J.; Zhang, C.; Cheng, J.; Wang, F.; Qiu, L. Effects of biogas residues containing antibiotics on soil enzyme activity and lettuce growth. *Environ. Sci. Pollut. Res.* **2019**, *26*, 6116–6122. [\[CrossRef\]](#) [\[PubMed\]](#)
31. Wojewódzki, P.; Lemanowicz, J.; Debska, B.; Haddad, S.A. Soil Enzyme Activity Response under the Amendment of Different Types of Biochar. *Agronomy* **2022**, *12*, 569. [\[CrossRef\]](#)
32. Cataldo, D.A.; Maroon, M.; Schrader, L.E.; Youngs, V.L. Rapid colorimetric determination of nitrate in plant tissue by nitration of salicylic acid. *Commun. Soil Sci. Plant Anal.* **1975**, *6*, 71–80. [\[CrossRef\]](#)
33. Baillie, I.C.; Anderson, J.M.; Ingram, J.S.I. Tropical Soil Biology and Fertility: A Handbook of Methods. *J. Ecol.* **1990**, *78*, 547. [\[CrossRef\]](#)
34. Bremner, J.M. Total Nitrogen. In *Methods of Soil Analysis, Part 2. Chemical and Microbiological Properties* (Agronomy 9); Norman, A.G., Ed.; American Society of Agronomy: Madison, WI, USA; Soil Science Society of America: Madison, WI, USA, 1965; pp. 1149–1178. [\[CrossRef\]](#)
35. Horneck, D.A.; Miller, R.O. Determination of total nitrogen in plant tissue. In *Handbook of Reference Methods for Plant Analysis*; Kalra, Y.P., Ed.; Soil and Plant Analysis Council, Inc.: Boca Raton, FL, USA; CRC Press: Boca Raton, FL, USA, 1998; pp. 75–83.
36. Blair, G.; Lefroy, R.; Lisle, L. Soil carbon fractions based on their degree of oxidation, and the development of a carbon management index for agricultural systems. *Aust. J. Agric. Res.* **1995**, *46*, 1459–1466. [\[CrossRef\]](#)
37. Garcia, C.; Hernandez, T.; Costa, F. Microbial activity in soils under mediterranean environmental conditions. *Soil Biol. Biochem.* **1994**, *26*, 1185–1191. [\[CrossRef\]](#)
38. Eivazi, F.; Tabatabai, M. Factors affecting glucosidase and galactosidase activities in soils. *Soil Biol. Biochem.* **1990**, *22*, 891–897. [\[CrossRef\]](#)
39. Biró, B.; Toscano, G.; Horváth, N.; Matics, H.; Domonkos, M.; Scotti, R.; Rao, M.A.; Wejden, B.; French, H.K. Vertical and horizontal distributions of microbial abundances and enzymatic activities in propylene-glycol-affected soils. *Environ. Sci. Pollut. Res.* **2014**, *21*, 9095–9108. [\[CrossRef\]](#) [\[PubMed\]](#)
40. Parham, J.; Deng, S. Detection, quantification and characterization of β -glucosaminidase activity in soil. *Soil Biol. Biochem.* **2000**, *32*, 1183–1190. [\[CrossRef\]](#)
41. Adam, G.; Duncan, H. Development of a sensitive and rapid method for the measurement of total microbial activity using fluorescein diacetate (FDA) in a range of soils. *Soil Biol. Biochem.* **2001**, *33*, 943–951. [\[CrossRef\]](#)
42. Watson, C.A.; Atkinson, D.; Gosling, P.; Jackson, L.R.; Rayns, F.W. Managing soil fertility in organic farming systems. *Soil Use Manag.* **2002**, *18*, 239–247. [\[CrossRef\]](#)
43. Uddin, S.; Islam, M.R.; Jahangir, M.M.R.; Rahman, M.M.; Hassan, S.; Hassan, M.M.; Abo-Shosha, A.A.; Ahmed, A.F.; Rahman, M.M. Nitrogen Release in Soils Amended with Different Organic and Inorganic Fertilizers under Contrasting Moisture Regimes: A Laboratory Incubation Study. *Agronomy* **2021**, *11*, 2163. [\[CrossRef\]](#)
44. Bengtsson, G.; Bengtson, P.; Månsson, K.F. Gross nitrogen mineralization-, immobilization-, and nitrification rates as a function of soil C/N ratio and microbial activity. *Soil Biol. Biochem.* **2003**, *35*, 143–154. [\[CrossRef\]](#)
45. Da Silva, A.; Rossetto, R.; Bonnetine, J.; Piemonte, M.; Muraoka, T. Net and Potential Nitrogen Mineralization in Soil with Sugarcane Vinasse. *Sugar Tech.* **2013**, *15*, 159–164. [\[CrossRef\]](#)
46. Argenta, G.; Da Silva, P.R.F.; Sangoi, L. Leaf relative chlorophyll content as an indicator parameter to predict nitrogen fertilization in maize. *Ciênc Rural* **2004**, *34*, 1379–1387. [\[CrossRef\]](#)
47. Lawlor, D.W.; Mengel, K.; Kirkby, E.A. Principles of plant nutrition. *Ann. Bot.* **2004**, *93*, 479–480. [\[CrossRef\]](#)
48. Maxwell, K.; Johnson, G.N. Chlorophyll fluorescence—A practical guide. *J. Exp. Bot.* **2000**, *51*, 659–668. [\[CrossRef\]](#) [\[PubMed\]](#)
49. Jägerbrand, A.K.; Kudo, G. Short-Term Responses in Maximum Quantum Yield of PSII (Fv/Fm) to ex situ Temperature Treatment of Populations of Bryophytes Originating from Different Sites in Hokkaido, Northern Japan. *Plants* **2016**, *5*, 22. [\[CrossRef\]](#) [\[PubMed\]](#)

50. Bjorkman, O.; Demmig, B. Photon yield of O₂ evolution and chlorophyll fluorescence characteristics at 77 K among vascular plants of diverse origins. *Planta* **1987**, *170*, 489–504. [\[CrossRef\]](#)
51. Bolhar-Nordenkamp, H.R.; Long, S.; Baker, N.R.; Oquist, G.; Schreiber, U.; Lechner, E.G. Chlorophyll Fluorescence as a Probe of the Photosynthetic Competence of Leaves in the Field: A Review of Current Instrumentation. *Funct. Ecol.* **1989**, *3*, 497. [\[CrossRef\]](#)
52. Maas, E.V.; Hoffman, G.J. Crop Salt Tolerance-Current Assessment. *J. Irrig. Drain. Div.* **1977**, *103*, 115–134. [\[CrossRef\]](#)
53. Wichern, J.; Wichern, F.; Joergensen, R.G. Impact of salinity on soil microbial communities and the decomposition of maize in acidic soils. *Geoderma* **2006**, *137*, 100–108. [\[CrossRef\]](#)
54. Iqbal, T.; Joergensen, R.G.; Knoblauch, C.; Lucassen, R.; Singh, Y.; Watson, C.; Wichern, F. Rice straw addition does not substantially alter microbial properties under hypersaline soil conditions. *Biol. Fertil. Soils* **2016**, *52*, 867–877. [\[CrossRef\]](#)
55. Chahal, S.S.; Choudhary, O.P.; Mavi, M.S. Organic amendments decomposability influences microbial activity in saline soils. *Arch. Agron. Soil Sci.* **2017**, *63*, 1875–1888. [\[CrossRef\]](#)
56. Leogrande, R.; Vitti, C. Use of organic amendments to reclaim saline and sodic soils: A review. *Arid Land Res. Manag.* **2018**, *33*, 1–21. [\[CrossRef\]](#)
57. Piccolo, A.; Pietramellara, G.; Mbagwu, J. Use of humic substances as soil conditioners to increase aggregate stability. *Geoderma* **1997**, *75*, 267–277. [\[CrossRef\]](#)
58. Zandonadi, D.B.; Santos, M.P.; Dobbss, L.B.; Olivares, F.L.; Canellas, L.P.; Binzel, M.L.; Okorokova-Façanha, A.L.; Façanha, A.R. Nitric oxide mediates humic acids-induced root development and plasma membrane H⁺-ATPase activation. *Planta* **2010**, *231*, 1025–1036. [\[CrossRef\]](#) [\[PubMed\]](#)
59. Havelcová, M.; Mizera, J.; Sýkorová, I.; Pekař, M. Sorption of metal ions on lignite and the derived humic substances. *J. Hazard. Mater.* **2009**, *161*, 559–564. [\[CrossRef\]](#) [\[PubMed\]](#)
60. Tang, W.-W.; Zeng, G.-M.; Gong, J.-L.; Liang, J.; Xu, P.; Zhang, C.; Huang, B.-B. Impact of humic/fulvic acid on the removal of heavy metals from aqueous solutions using nanomaterials: A review. *Sci. Total. Environ.* **2014**, *468–469*, 1014–1027. [\[CrossRef\]](#) [\[PubMed\]](#)
61. Yao, J.; Kong, Q.; Qiu, Z.; Chen, L.; Shen, D. Patterns of heavy metal immobilization by MSW during the landfill process. *Chem. Eng. J.* **2019**, *375*, 122060. [\[CrossRef\]](#)
62. Masciandaro, G.; Ceccanti, B. Assessing soil quality in different agro-ecosystems through biochemical and chemico-structural properties of humic substances. *Soil Tillage Res.* **1999**, *51*, 129–137. [\[CrossRef\]](#)
63. Gramss, G.; Ziegenhagen, D.; Sorge, S. Degradation of Soil Humic Extract by Wood- and Soil-Associated Fungi, Bacteria, and Commercial Enzymes. *Microb. Ecol.* **1999**, *37*, 140–151. [\[CrossRef\]](#)
64. García, A.C.; Santos, L.A.; Izquierdo, F.G.; Sperandio, M.V.L.; Castro, R.N.; Berbara, R.L.L. Vermicompost humic acids as an ecological pathway to protect rice plant against oxidative stress. *Ecol. Eng.* **2012**, *47*, 203–208. [\[CrossRef\]](#)
65. Killham, K.; Firestone, M.K. Proline transport increases growth efficiency in salt-stressed *Streptomyces griseus*. *Appl. Environ. Microbiol.* **1984**, *48*, 239–241. [\[CrossRef\]](#)
66. Schimel, J.P.; Scott, W.J.; Killham, K. Changes in Cytoplasmic Carbon and Nitrogen Pools in a Soil Bacterium and a Fungus in Response to Salt Stress. *Appl. Environ. Microbiol.* **1989**, *55*, 1635–1637. [\[CrossRef\]](#)
67. European Commission. Commission Regulation (EU) No 1258/2011 of 2 December 2011 amending Regulation (EC) No. 1881/2006 as regards maximum levels for nitrates in foodstuffs. *Off. J. Eur. Union* **2011**, *L320*, 15–17.
68. Ayyobi, H.; Olfati, J.-A.; Peyvast, G.-A. The effects of cow manure vermicompost and municipal solid waste compost on peppermint (*Mentha piperita* L.) in Torbat-e-Jam and Rasht regions of Iran. *Int. J. Recycl. Org. Waste Agric.* **2014**, *3*, 147–153. [\[CrossRef\]](#)
69. Mills, H.A.; Jones, J.B., Jr. *Plant Analysis Handbook II. A Practical Sampling, Preparation, Analysis, and Interpretation Guide*; Micro-Macro Publishing: Athens, Greece, 1996.
70. Zahidah, A.R.; Rosenani, A.B.; Hajar, A.S.; Nozulaidi, N.M. Effects of Different Types of Organic Fertilizer on Biomass Yield, Bioactive Compounds and Heavy Metals Contents of *Phyllanthus Niruri*. *J. Phys. Conf. Ser.* **2021**, *2000*, 012005. [\[CrossRef\]](#)
71. Wang, L.; Liu, S.; Li, J.; Li, S. Effects of Several Organic Fertilizers on Heavy Metal Passivation in Cd-Contaminated Gray-Purple Soil. *Front. Environ. Sci.* **2022**, *10*, 884. [\[CrossRef\]](#)
72. Wikar, J. The effect of fertilizing soils degraded by the metallurgical industry on the content of elements in *Lactuca sativa* L. *Sci. Rep.* **2021**, *11*, 4072. [\[CrossRef\]](#)
73. Yang, F.; Tang, C.; Antonietti, M. Natural and artificial humic substances to manage minerals, ions, water, and soil microorganisms. *Chem. Soc. Rev.* **2021**, *50*, 6221–6239. [\[CrossRef\]](#) [\[PubMed\]](#)
74. Zeng, F.; Ali, S.; Zhang, H.; Ouyang, Y.; Qiu, B.; Wu, F.; Zhang, G. The influence of pH and organic matter content in paddy soil on heavy metal availability and their uptake by rice plants. *Environ. Pollut.* **2011**, *159*, 84–91. [\[CrossRef\]](#) [\[PubMed\]](#)
75. Han, H.; Wu, X.; Bolan, N.; Kirkham, M.; Yang, J.; Chen, Z. Inhibition of cadmium uptake by wheat with urease-producing bacteria combined with sheep manure under field conditions. *Chemosphere* **2022**, *293*, 133534. [\[CrossRef\]](#)
76. Demisie, W.; Liu, Z.; Zhang, M. Effect of biochar on carbon fractions and enzyme activity of red soil. *Catena* **2014**, *121*, 214–221. [\[CrossRef\]](#)
77. Dalal, R. Urease activity in some Trinidad soils. *Soil Biol. Biochem.* **1975**, *7*, 5–8. [\[CrossRef\]](#)
78. Jha, D.K.; Sharma, G.D.; Mishra, R.R. Soil microbial population numbers and enzyme activities in relation to altitude and forest degradation. *Soil Biol. Biochem.* **1992**, *24*, 761–767. [\[CrossRef\]](#)

79. Amador, J.A.; Glucksman, A.M.; Lyons, J.B.; Gorres, J.H. Spatial distribution of soil phosphatase activity within a riparian forest. *Soil Sci.* **1997**, *162*, 808–825. [\[CrossRef\]](#)
80. Tabatabai, M.A. Soil enzymes. In *Methods of Soil Analysis, Part 2 Microbiological and Biochemical Properties*; Mickelson, S.H., Bighan, J.M., Eds.; Soil Science Society of America: Madison, WI, USA, 1994; pp. 775–826.
81. Bandick, A.K.; Dick, R.P. Field management effects on soil enzyme activities. *Soil Biol. Biochem.* **1999**, *31*, 1471–1479. [\[CrossRef\]](#)
82. Kızılkaya, R.; Bayraklı, B. Effects of N-enriched sewage sludge on soil enzyme activities. *Appl. Soil Ecol.* **2005**, *30*, 192–202. [\[CrossRef\]](#)
83. Gu, Y.; Wang, P.; Kong, C. Urease, invertase, dehydrogenase and polyphenoloxidase activities in paddy soil influenced by allelopathic rice variety. *Eur. J. Soil Biol.* **2009**, *45*, 436–441. [\[CrossRef\]](#)
84. Ajwa, H.A.; Tabatabai, M.A. Decomposition of different organic materials in soils. *Biol. Fertil. Soils* **1994**, *18*, 175–182. [\[CrossRef\]](#)
85. Turner, B.L.; Hopkins, D.W.; Haygarth, P.M.; Ostle, N. β -Glucosidase activity in pasture soils. *Appl. Soil Ecol.* **2002**, *20*, 157–162. [\[CrossRef\]](#)
86. Filipek-Mazur, B.; Pużyńska, K.; Tabak, M.; Pużyński, S. Enzymatic Activity of Soil under Spelt Grown in an Organic Farming System in Poland's Temperate Climate. *Agronomy* **2020**, *10*, 930. [\[CrossRef\]](#)
87. Wang, H.; Boutton, T.; Xu, W.; Hu, G.; Jiang, P.; Bai, E. Quality of fresh organic matter affects priming of soil organic matter and substrate utilization patterns of microbes. *Sci. Rep.* **2015**, *5*, 10102. [\[CrossRef\]](#)
88. Chathurika, J.A.S.; Kumaragamage, D.; Indraratne, S.P.; Dandeniya, W.S. Improving soil carbon pool, soil fertility and yield of maize (*Zea mays* L.) in low-fertile tropical Alfisols by combining fertilizers with slow-decomposing organic amendments. *J. Agric. Sci.* **2019**, *157*, 45–54. [\[CrossRef\]](#)

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.