

Infection, Genetics and Evolution

Investigating the genetic structure of the parasites *Anisakis pegreffii* and *A. berlandi* (Nematoda: Anisakidae) in a sympatric area of the southern Pacific Ocean waters using a multilocus genotyping approach: first evidence of their interspecific hybridization

--Manuscript Draft--

Manuscript Number:	
Article Type:	VSI: Helminths
Keywords:	Anisakis, New Zealand, SSR loci, nas10 nDNA, mtDNA cox2, sex-linked loci, hybridization
Corresponding Author:	Simonetta Mattiucci, PhD Sapienza University of Rome Rome, ITALY
First Author:	Eleonora Bello
Order of Authors:	Eleonora Bello Marialetizia Palomba Stephen C. Webb Michela Paoletti Paolo Cipriani Giuseppe Nascetti Simonetta Mattiucci, PhD
Abstract:	The southern Pacific Ocean, off the New Zealand coast, has been reported as one sympatric area of the two parasite species <i>Anisakis pegreffii</i> and <i>A. berlandi</i>. Here, a multilocus genotyping approach, based on a panel of eleven DNA microsatellite (SSR) loci plus the sequences analysis of the nuclear nas10 nDNA and the mitochondrial mtDNA cox 2 gene loci, was applied to a total of N= 344 adults and larvae of <i>Anisakis</i> spp. from cetacean and fish species, respectively. Out of the newly scored SSR loci, Anisl 15 and Anisl 2 showed fixed alternative alleles between <i>A. pegreffii</i> and <i>A. berlandi</i> resulting as 100% diagnostic loci. Out of SSRs Anisl 00314 and Anisl 7 previously disclosed, two additional loci, i.e., Anisl 4 and Anisl 22, were found to be sex-linked. The Bayesian genotypes clustering approach (STRUCTURE) allowed identification, with a 100% of probability value, N= 208 specimens to the "pure parental" <i>A. pegreffii</i>, N= 133 to the "pure parental" <i>A. berlandi</i>, while one adult and two larval stages showed mixed ancestry between the two groups having, in all cases, a Q-value= 0.50. NEWHYBRIDS analysis assigned (100% of probability) those specimens to their F1 hybrid category. This represents the first evidence of contemporary hybridization between the two parasite species in a sympatric area. The pairwise F_{ST} values estimated at intraspecific and interspecific level, inferred from both SSR loci and mitochondrial mtDNA cox 2 sequences, have also demonstrated the existence of two distinct panmictic units in this study area, corresponding respectively to <i>A. pegreffii</i> and <i>A. berlandi</i>. The results obtained support the useful application of a multilocus approach in the identification of sibling species and their hybrid categories in sympatric areas. The possible use of sex-linked SSR loci of the two species of the <i>A. simplex</i> (s. l.), for sex determination of their larval stages, is also suggested.
Suggested Reviewers:	Elżbieta Łopieńska-Biernat associate professor ela.lopienska@uwm.edu.pl Shokoofeh Shamsi associate professor sshamsi@csu.edu.au Melanie Gay

	melanie.GAY@anses.fr
	Regina Klapper regina.klapper@mri.bund.de

DIPARTIMENTO DI SANITÀ
PUBBLICA E MALATTIE INFETTIVE



SAPIENZA
UNIVERSITÀ DI ROMA

Simonetta Mattiucci, PhD
Associate Professor
Department of Public Health and Infectious Diseases
Head of "Parasitology Unit" at the Academic Hospital "Policlinico Umberto I"
"Sapienza - University of Rome"
P.le Aldo Moro, 5 00185
Rome Italy
Tel ++39 06 49914894
email: simonetta.mattiucci@uniroma1.it

Prof. Michel Tibayrenc
Editor-in-Chief of
Infection, Genetics and Evolution

Dear Professor Michel Tibayrenc,

please find, here enclosed, the file of the MS entitled: "*Investigating the genetic structure of the parasites Anisakis pegreffii and A. berlandi (Nematoda: Anisakidae) in a sympatric area of the southern Pacific Ocean waters using a multilocus genotyping approach: first evidence of their interspecific hybridization*" by: Eleonora Bello, Marialetizia Palomba, Stephen C. Webb, Michela Paoletti, Paolo Cipriani, Giuseppe Nascetti, and Simonetta Mattiucci for its submission to the Special Issue "*Helminths*" of Your Journal. The MS text includes 6 Tables and 4 Figures (plus 1 Suppl. Fig).

All the Authors have read the MS; they agree with the content of this version of the MS.

The Authors declare that they have no conflicting interests.

The MS is not currently submitted for publication elsewhere.

We do hope that you wish to consider the MS as suitable for its publication on Your Journal.

Waiting for your reply,

with kind regards,

Simonetta Mattiucci

Rome, 05.03.2021

1 **Highlights**

- 2 1. Multilocus genotyping in the genetic structure of *A. pegreffii* and *A. berlandi*
- 3 2. Four SSR loci were newly scored in the two parasite species
- 4 3. *Anisl* 2 and *Anisl* 15 are 100% diagnostic between the two species
- 5 4. Two additional SSR loci were disclosed as sex-linked in the two species
- 6 5. First evidence of contemporary hybridization between the two species

Investigating the genetic structure of the parasites *Anisakis pegreffii* and *A. berlandi* (Nematoda: Anisakidae) in a sympatric area of the southern Pacific Ocean waters using a multilocus genotyping approach: first evidence of their interspecific hybridization

Eleonora Bello^{a,b,§}, Marialetizia Palomba^{b,c,§}, Stephen C. Webb^d, Michela Paoletti^a, Paolo Cipriani^e, Giuseppe Nascetti^a, Simonetta Mattiucci^{b*}

^a*Department of Biological and Ecological Sciences, Tuscia University, Viale dell'Università snc, 01100 Viterbo - Italy*

^b*Department of Public Health and Infectious Diseases, Section of Parasitology, Sapienza University of Rome. P.le Aldo Moro 5, 00185 Rome - Italy; Laboratory affiliated to "Istituto Pasteur Italia - Fondazione Cenci-Bolognetti"*

^c*Department of Integrative Marine Ecology, Stazione Zoologica Anton Dohrn. Villa Comunale, 80121 Naples - Italy*

^d*Cawthron Institute, Nelson, New Zealand*

^e*Institute of Marine Research (IMR), Bergen, Norway*

§ equal contribution

*Corresponding author: Department of Public Health and Infectious Diseases, Section of Parasitology, Sapienza - University of Rome, Italy, P.le Aldo Moro, 5 00189 Rome, Italy
Laboratory affiliated to "Istituto Pasteur Italia - Fondazione Cenci Bolognetti"

E-mail address: simonetta.mattiucci@uniroma1.it

Abstract

The southern Pacific Ocean, off the New Zealand coast, has been reported as one sympatric area of the two parasite species *Anisakis pegreffii* and *A. berlandi*. Here, a multilocus genotyping approach, based on a panel of eleven DNA microsatellite (SSR) loci *plus* the sequences analysis of the nuclear *nas10* nDNA and the mitochondrial mtDNA *cox2* gene loci, was applied to a total of N= 344 adults and larvae of *Anisakis* spp. from cetacean and fish species, respectively. Out of the newly scored SSR loci, *Anisl* 15 and *Anisl* 2 showed fixed alternative alleles between *A. pegreffii* and *A. berlandi* resulting as 100% diagnostic loci. Out of SSRs *Anisl* 00314 and *Anisl* 7 previously disclosed, two additional loci, i.e., *Anisl* 4 and *Anisl* 22, were found to be sex-linked. The Bayesian genotypes clustering approach (STRUCTURE) allowed identification, with a 100% of probability value, N= 208 specimens to the "pure parental" *A. pegreffii*, N= 133 to the "pure parental" *A. berlandi*, while one adult and two larval stages showed mixed ancestry between the two groups having, in all cases, a Q-value= 0.50. NEWHYBRIDS analysis assigned (100% of probability) those specimens to their F1 hybrid category. This represents the first evidence of contemporary hybridization between the two parasite species in a sympatric area. The pairwise F_{ST} values estimated at intraspecific and interspecific level, inferred from both SSR loci and mitochondrial mtDNA *cox2* sequences, have also demonstrated the existence of two distinct panmictic units in this study area, corresponding respectively to *A. pegreffii* and *A. berlandi*. The results obtained support the useful application of a multilocus approach in the identification of sibling species and their hybrid categories in sympatric areas. The possible use of sex-linked SSR loci of the two species of the *A. simplex* (s. l.), for sex determination of their larval stages, is also suggested.

Keywords: *Anisakis*, New Zealand, SSR loci, *nas10* nDNA, mtDNA *cox2*, sex-linked loci, hybridization

50
1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1. Introduction

Nematode parasites of the genus *Anisakis* have a heteroxenous life cycle which involves various marine organisms at different levels of the trophic web in the marine ecosystem. The adults live in the stomachs of marine mammals, mainly cetaceans (definitive hosts); fish and squid are intermediate or paratenic hosts, while planktonic or semi-planktonic crustaceans act as first intermediate hosts. To date, nine nominal species and two undescribed genotypes are recognised in the genus *Anisakis*, as based on genetic/molecular systematics (for a review, Mattiucci et al., 2018a). Indeed, the existence of nine species as distinct phylogenetic units has been demonstrated by various concatenated phylogenetic analyses, as inferred by combining both nuclear and mitochondrial sequences data (Valentini et al., 2006; Mattiucci and Nascetti, 2008; Cavallero et al., 2011; Mattiucci et al., 2014, 2018a). According to those analyses, four distinct clades within the genus *Anisakis* exist (Mattiucci et al., 2018a). Among them, one clade encompasses the three sibling species of the *A. simplex* (s. l.) complex: *A. simplex* (s. s.), *A. pegreffii* and *A. berlandi* (Nascetti et al., 1986; Mattiucci et al., 2014). Genetic/molecular markers mostly used for the recognition of the three sibling species of the *A. simplex* (s. l.) complex were inferred from mitochondrial gene loci (the direct sequences analysis of the mtDNA *cox2*) and nuclear loci, such as the allozymes (Mattiucci et al., 1997, 2016), the direct sequences analysis and the RFLPs-PCR of the ITS region of rDNA (D'Amelio et al., 2000), and the EF-1-alpha nDNA (Mattiucci et al., 2016); the last gene locus, however, was developed only in the species *A. pegreffii* and *A. simplex* (s. s.). More recently, the existence of those three species as distinct biological species, reproductively isolated, has been supported also using polymorphic DNA microsatellite loci (SSRs) (Mattiucci et al., 2019; Bello et al., 2020) and direct sequencing of the *nas10* nDNA (Palomba et al., 2020b).

Anisakis pegreffii has been so far recorded as a parasite, at the adult stage, of mainly oceanic dolphins belonging to the Families Delphinidae, Monodontidae and Phocoenidae and, less

frequently, in Neobalenidae whales (Mattiucci et al., 2018a). At its third larval stage, it occurs in more than 60 pelagic and benthopelagic fish species, and in four squid species throughout its geographical range (Mattiucci et al., 2018a). *Anisakis pegreffii* is the most common species of *Anisakis* in the Mediterranean Sea, while in the Atlantic waters the northerly limit of its geographical range is represented by the Iberian coast, from where the species has been frequently identified in sympatry with *A. simplex* (s. s.) in several demersal and pelagic fish species and as well as adults in cetaceans. However, *A. pegreffii* is also widespread in the Pacific Boreal region (Japan Sea, China Sea) and in the Austral Region, between 30°S and 60°S, where the species was identified at both adult and larval stages (reviewed in Mattiucci et al., 2018a; Gomes et al., 2020, 2021).

The species *A. berlandi* (see Mattiucci et al., 2014) currently exhibits a discontinuous distribution, being reported in the Southern Region, as in the Chilean Pacific Ocean, the South Shetland Islands, New Zealand waters and the South African Atlantic coast (Mattiucci and Nascetti, 2008; Klimpel et al., 2010; Mattiucci et al., 2018a). From these geographical areas, *A. berlandi* has been identified at the adult stage in six cetacean species, while its Type I larvae were so far identified in fish species from Southern waters from off New Zealand (Mattiucci et al., 2014; 2019; Bello et al., 2020), as well as from Southern Chilean (Mattiucci et al., 2018a) and Argentine (Irigoitia et al., 2018) coasts. Fourth-stage larvae of *A. berlandi* were also identified in *Kogia sima* from Australian waters (Shamsi et al., 2012) and, as L3, in other accidental hosts of the New Caledonian waters (Shamsi et al., 2017). Klimpel et al. (2010) stated that the occurrence of few larval specimens of *A. berlandi* and *A. pegreffii* in myctophids from the southern waters of the South Shetland Islands (Antarctic area) could be related to the introduction of those two parasites species from outside the Antarctic, through their migrating fish intermediate hosts. Accordingly, a very low occurrence of *A. berlandi* was recorded also in *M. leonina* from the South Shetland Islands (Mattiucci and Nascetti, 2008).

So far, the sympatric occurrence of *A. pegreffii* and *A. berlandi* is recorded in the South Pacific (i.e., in the New Zealand waters) (Mattiucci et al., 1997; 2014) and in the South Atlantic Ocean (Argentine coast) (Irigoitia et al., 2018), in Chile and South Africa (Mattiucci et al., 2018a). However, although their range of distribution overlap in some geographic areas, the evidence of hybridization events between *A. pegreffii* and *A. berlandi* has not been so far recorded. It has been also indicated that nuclear multilocus genotyping approach should be used for investigating events such as hybridization, introgression and the retention of ancestral polymorphism in closely related anisakid species in sympatric areas (Mattiucci et al., 2016; 2018a). Thus, in recent years, in order to recognise parental genotypes of the species of the *A. simplex* (s. l.) complex and their mixed ancestry in known overlapping ranges of distribution, novel diagnostic markers from the nuclear genomes of these parasites have been developed (Mattiucci et al., 2016; 2019; Gomez-Mateos et al., 2020; Bello et al., 2020; Palomba et al., 2020b).

In the present paper, a multilocus genotyping approach, inferred from the screening of a panel of 11 microsatellite DNA loci (SSRs) and the sequences analysis of the *nas10* nDNA and the mtDNA *cox2* gene loci, was carried out on adult and larval populations of *A. pegreffii* with populations of the sibling species *A. berlandi* detected in sympatry from one of their contact areas (i.e. New Zealand waters). The aim of this work is to: i) assign the specimens to the two parental taxa, i.e. *A. pegreffii* or *A. berlandi*; ii) recognise their mixed ancestry based on a Bayesian multinuclear clustering approach; and iii) infer data on the genetic diversity, at both nuclear and mitochondrial level, of the two species from this study area.

2. Materials and Methods

2.1 Collection of parasite samples

A total of $N= 225$ adults (A) and fourth-stage larvae (L4) of nematodes from four cetacean species referred to the genus *Anisakis*, and $N= 119$ third-stage (L3) *Anisakis* Type I larvae (*sensu* Berland, 1961), collected from eight fish species from the South Pacific waters of the New Zealand coast, were genotyped by different genetic markers. Details concerning the life-history stage, definitive hosts (cetacean species) and intermediate/paratenic (fish species) of the *Anisakis* spp. specimens examined are given in Table 1. In particular, adult nematodes were from cetaceans stranded on the coast of New Zealand, i.e., three pilot whales, *Globicephala melas* (Traill), one Risso's dolphin, *Grampus griseus* (Cuvier), a Hector's dolphin, *Cephalorhynchus hectori* (van Beneden), and a common dolphin *Delphinus delphis* Linnaeus, 1758. The L3-stage larvae were collected from different fish species (Table 1) caught during the winter in 2018 along the southern coast of New Zealand.

The nematodes were repeatedly washed in saline solution and then preserved in 70% alcohol. Considering the sex-linkage of microsatellite loci, as discovered in our previous studies (Mattiucci et al., 2019; Bello et al., 2020), the adult specimens were first distinguished as females and males; in particular, sex discrimination was performed according to the main morphological diagnostic features between sexes (Mattiucci et al., 2014), using an optical microscope, at a total magnification of 100–400X. The numbers of adult female and male specimens studied are reported in Table 1. In the case of adult female worms, a tissue sample for molecular analysis was taken as distant as possible from the uterus to avoid possible contamination from stored male's sperm.

2.2 Genetic analysis of *Anisakis* spp. specimens by sequencing of *nas10* nDNA and mtDNA *cox2*

All ($N= 344$) the specimens here studied were first assigned to their species level by the sequences analysis of the recently discovered as nuclear gene locus that is diagnostic between

the three species of the *A. simplex* (s. l.) complex, i.e., the *nas10* nDNA (Palomba et al., 2020b). Additionally, the same larval and adult specimens ($N= 344$) were also sequenced at the mtDNA *cox2* gene locus, which is a standardised marker to recognise species of the genus *Anisakis* (Mattiucci et al., 2014).

Total DNA was extracted from a tissue portion (~2 mg) of each larva and adult specimen of *Anisakis* (Table 1). The extraction method by Quick-gDNA Miniprep Kit (ZYMO RESEARCH) was used. DNA obtained was quantified using a NanoDrop_TC1-E20 spectrophotometer (BioTek Synergy HT).

For the direct sequencing of the *nas10* nDNA gene, the PCR amplification was performed according to the procedures detailed in Palomba et al. (2020b), using the primers *nas10-F* (5'-GAT GTT CCT GCA AGT GAT TG -3') and *nas10-R* (5'- CGC TAT TAA GAG AGG GAT CG - 3').

For the sequencing of the mtDNA *cox2* gene locus, PCR amplification was performed using the primers 211F (5'-TTT TCT AGT TAT ATA GAT TGR TTT YAT-3') and 210R (5'-CAC CAA CTC TTA AAA TTA TC-3'). PCR conditions were established as previously described (Mattiucci et al., 2014).

2.3 PCR amplification of SSR markers and genotyping in *Anisakis pegreffii* and *A. berlandi*

The same specimens ($N= 344$) of *A. pegreffii* and *A. berlandi*, were also tested at the panel of seven SSR loci among those previously developed by Mattiucci et al., 2019 (i.e., *Anisl 00185*, *Anisl 00314*, *Anisl 10535*, *Anisl 05784*, *Anisl 08059* and *Anisl 00875*) and Mladineo et al., 2017 (i.e., *Anisl 7*) for the species *A. pegreffii* and *A. simplex* (s. s.) (Mattiucci et al., 2019), as well as in *A. berlandi* (Bello et al., 2020). Additionally, a further four SSR loci (named as *Anisl 4*, *Anisl*

22, *Anisl 15* and *Anisl 2*), among those developed in Mladineo et al. (2017), were here tested for the first time on the collected specimens (Table 1).

The amplification of microsatellite loci was performed in three Multiplex PCRs: *Anisl 05784*, *Anisl 08059* and *Anisl 00875* in Multiplex 1; *Anisl 00185*, *Anisl 00314*, *Anisl 10535* and *Anisl 7* in Multiplex 2; and *Anisl 4*, *Anisl 22*, *Anisl 15* and *Anisl 2* in Multiplex 3. Each Multiplex PCR amplification was performed in a 10 µl reaction volume, containing: 5-10 ng of genomic DNA, 5 µl Type-it Microsatellite PCR Kit (Qiagen®), double distilled water, and concentrations of 10 µM labeled forward and reverse primers each (Mattiucci et al., 2019). The following cycling protocol was used for the amplification for the three multiplex reactions: 35 cycles with 94°C for 30 sec, 56°C for 90 sec and 72°C for 60 sec. Before the first cycle, a prolonged denaturation step (95°C for 15 min) was included and the last cycle was followed by a 15 min extension at 60°C (Mattiucci et al., 2019).

Amplified PCR products were genotyped by an external Company (Macrogen service). The software Genemapper v.4.1 (Applied Biosystems, USA) was used to analyse the alleles obtained from the electropherograms. Genotyping errors generally associated with microsatellite analysis, such as stutter bands, the presence of null alleles and allelic drop-out were verified by using the software MICRO-CHECKER version 2.2.3 (Van Oosterhout et al., 2004).

2.4 Genetic data analysis

The sequences obtained at the *nas10* nDNA and the mtDNA *cox2* gene regions from the N=344 studied nematodes were aligned by using Clustal X (version 2.0) software (Larkin et al., 2007) with those of the same species previously obtained and deposited in GenBank (Mattiucci et al., 2014; Palomba et al., 2020b). This allowed the detection of those fixed diagnostic nucleotide positions that discriminate the species *A. pegreffii* and *A. berlandi*.

The observed heterozygosity (H_o), the expected heterozygosity (H_e), the mean number of alleles per locus (A) observed at the 11 SSR loci were calculated by ARLEQUIN version 3.5 (Excoffier and Lischer, 2010). The occurrence of the expected Hardy–Weinberg Equilibrium (HWE) at each SSR locus was assessed by means of exact tests, as implemented in the software ARLEQUIN version 3.5 (Excoffier and Lischer, 2010). Analysis of molecular variance (AMOVA), as implemented in ARLEQUIN version 3.5, with 1023 permutations, was applied on the genetic data sets obtained in the populations of *A. pegreffii* and *A. berlandi*. The fixation index (F_{IS}) from the genetic data sets obtained at the SSR loci in the analysed populations of the two *Anisakis* species, was estimated using ARLEQUIN version 3.5 (Excoffier and Lischer, 2010). The F_{ST} (Weir and Cockerham, 1984) values, inferred from both SSR loci and mtDNA *cox2* genetic data sets, were estimated by ARLEQUIN version 3.5 (Excoffier and Lischer, 2010).

To investigate instances of gene exchange between the two species, a Bayesian clustering algorithm was used from the SSRs and the *nas10* nDNA sequences data sets, performed by the program STRUCTURE v. 2.3.3 (Pritchard et al., 2000). The analysis was run setting the predefined number of clusters between 1 and 3. Twenty replicates of the analysis were carried out to check for consistency, each run for 100 000 MCMC iterations, following a burn-in of 50 000 iterations, under the admixture model and the assumption of correlated allele frequencies among populations. The best K value was identified using both the log probability of the data and the rate of change in the log probability of the data between successive K values as the optimality criteria (Evanno et al., 2005). The Bayesian analysis implemented in NEWHYBRIDS v. 1.1 (Anderson and Thompson, 2002) was also used to assess the occurrence of hybridization events between *A. pegreffii* and *A. berlandi*. This analysis assigns, probabilistically, each sampled individual to a predefined hybrid class, that is: pure parental (*A. pegreffii* or *A. berlandi*), first generation hybrid (F1), second generation hybrid (F2, or backcrosses). The assignment was performed on the SSRs and *nas10* nDNA data sets by computing 100 000

MCMC iterations, discarding the first 20000 iterations as burn-in (after checking for stationarity), and replicating the analysis 10 times to check for consistency.

3. Results

3.1 *Anisakis pegreffii* and *A. berlandi* identification based on *nas10* nDNA and mtDNA *cox2* sequences analysis

A total of $N = 344$ larval and adult specimens of *Anisakis* spp. (Table 1) were identified by sequences analysis of the nuclear *nas10* nDNA region. A fragment of 451 bp of the *nas10* nDNA region was obtained for all 344 specimens analysed. It revealed the presence of the nucleotide site diagnostic between the two species, as previously indicated in Palomba et al. (2020a). Indeed, at the position 251, a total of $N = 208$ individuals always showed the homozygote genotype G/G, while $N = 133$ specimens were always C/C homozygous (Fig. 1A). According to this result, 208 specimens were assigned to the parental *A. pegreffii*, whereas 133 were assigned to the parental species *A. berlandi* (s. s.) (Table 1). Patterns of heterozygote genotypes (G/C), i.e., showing an overlapping peak at that diagnostic position (Fig. 1B), were observed in three specimens (one adult collected from *G. melas* and two L3 larvae from *T. declivis*) (Table 2).

Sequences of the *nas10* nDNA region were deposited in GenBank under the accession numbers from MW691969 to MW691972 for *A. pegreffii*, and from MW691973 to MW691976, for *A. berlandi*.

The obtained sequences at the mtDNA *cox2* locus (629 bp) in 208 specimens matched at the 99% or 100% the sequences *A. pegreffii* previously deposited by us in GenBank (Table 2), while $N = 136$ were assigned to *A. berlandi* species (Table 2). Indeed, those 2 larval specimens from the fish host *T. declivis* and 1 female adult specimen from *G. melas*, which have shown a heterozygote genotype at the *nas10* nDNA, according to the mtDNA *cox2* sequences resulted as belonging to *A. berlandi* (Table 2).

Sequences of the mtDNA *cox2* region were deposited in GenBank under the accession numbers from MW658562 to MW658565 for *A. pegreffii*, and from MW658566 to MW658569, for *A. berlandi*.

3.2 Genetic diversity of SSR loci in *A. pegreffii* and *A. berlandi*

The same individual specimens ($N= 344$) previously sequenced at the nuclear gene *nas10* nDNA and mtDNA *cox2* gene regions, once genotyped at the eleven scored SSR loci, have shown some differences in the overall genetic diversity values in the two species, based on both the mean number of alleles per locus, and values of expected heterozygosity per locus (H_e) (Table 3).

Six loci (i.e., *Anisl 00185*, *Anisl 10535*, *Anisl 05784*, *Anisl 08059*, *Anisl 00875* and *Anisl 7*) were polymorphic in *A. pegreffii* and *A. berlandi*. Analogously, the newly tested SSR locus *Anisl 4* also was polymorphic in both parasite species. In contrast, the loci *Anisl 15* and *Anisl 2*, also here newly scored for the two *Anisakis* species, were found to be monomorphic. Additionally, in *A. berlandi*, only one allele was observed at the loci *Anisl 00314* and in the new tested *Anisl 22*; in contrast, the last locus showed 8 distinct alleles in *A. pegreffii* (Table 3).

The total number of alleles in *A. pegreffii* varied between 4, such as those observed at the locus *Anisl 7* in both larvae and adults, and the 14 found at the polymorphic locus *Anisl 08059* (Table 3). In the larval and adult populations of *A. pegreffii*, the mean value of alleles per locus (A), based on 11 loci, resulted as $A \approx 8.0$. In *A. berlandi*, the total number of alleles at the same loci was found to vary between 2, found in the loci *Anisl 10535* and *Anisl 05784*, up to 21 observed in the adult and larval specimens at the locus *Anisl 7* (Table 3). In *A. berlandi* the mean value of alleles per locus (A) was $A \approx 5.45$.

Deviation from the Hardy-Weinberg Equilibrium (HWE) at each locus was tested in the adult and larval populations of the two species (Table 3). No significant deviation between observed

(*Ho*) and expected (*He*) heterozygosity was seen at the loci *Anisl* 00185, *Anisl* 10535, *Anisl* 05784, *Anisl* 08059 and *Anisl* 00875 in the analysed samples of *A. pegreffii* and *A. berlandi* (Table 3). In contrast, statistically high significant departures from the HWE in both larval and adult populations of *A. berlandi* were observed at the locus *Anisl* 7, and in the newly scored *Anisl* 4. Analogously, both adult and larval populations of *A. pegreffii* showed a significant deviation of *Ho* from the *He* at the newly scored loci *Anisl* 4 and *Anisl* 22, as well as at the *Anisl* 00314 and *Anisl* 7 (Table 3).

3.3 New sex-linked SSR loci in *A. pegreffii* and *A. berlandi*

A positive F_{IS} value, indicating a marked excess of homozygotes from the expected HWE, was found at some SSR loci (Fig. 2A). In particular, significant positive F_{IS} values were recorded in *Anisl* 00314, *Anisl* 7 and in the newly studied loci, *Anisl* 4 and *Anisl* 22 (Fig. 2A). However, when considering only adult specimens and dividing the genotypes observed into adult male and female worms at the loci *Anisl* 00314, *Anisl* 7, *Anisl* 4 and *Anisl* 22 in *A. pegreffii*, as well as *Anisl* 7 and *Anisl* 4 in *A. berlandi*, the F_{IS} value in the male worms of the two species was equal to 1, showing a homozygous genotype (Fig. 2B). Therefore, out of the loci *Anisl* 00314 and *Anisl* 7, already known to be sex-linked loci in both species, also the loci *Anisl* 4 and *Anisl* 22 were seen to be sex-linked because of the hemizyosity state of the males in both the *Anisakis* spp., as X-linked inheritance. Indeed, the HWE again calculated at sex-linked loci considering only adult females of the two species, i.e., $N=71$ of *A. pegreffii* and $N=22$ of *A. berlandi*, showed no significant deviations from the HWE expectations (Table 4).

3.4 Multi-locus genotyping approach to distinguishing the two parental species and their hybrids

Allele frequencies estimated in *A. pegreffii* and *A. berlandi* are reported in Table 5. As regards the sex-linked loci (i.e., *Anisl* 00314, *Anisl* 7, *Anisl* 4 and *Anisl* 22), the allele frequencies were calculated considering the hemizygosity of male specimens, according to Mattiucci et al. (2019): male specimens were considered as monoallelic, while their female counterparts were considered as biallelic. Among the 11 SSR loci considered, *Anisl* 10535, *Anisl* 05784, *Anisl* 00875, *Anisl* 7 and *Anisl* 22 have shown a differential frequency distribution at some alleles found in common in the two species (Table 5). Instead, at the loci *Anisl* 15 and *Anisl* 2, *N*= 208 specimens of *A. pegreffii* were monomorphic, showing fixed alternative alleles (Table 5) with respect to those alleles observed in the specimens (*N*= 133) of *A. berlandi*. Indeed, the loci *Anisl* 15 and *Anisl* 2 were 100% diagnostic between the two parasite species (Fig. 3).

Heterozygote genotypes formed by those two fixed alternative alleles at the diagnostic loci *Anisl* 15 and *Anisl* 2 were observed in three specimens (Fig. 3): they were the same individual worms (Table 2) which have shown also a heterozygous genotype (C/G) at the diagnostic nucleotide site of *nas10* nDNA locus between the two parasite species (Fig. 1B); those three individuals, at the other partially diagnostic SSR loci (i.e., *Anisl* 10535, *Anisl* 05784, *Anisl* 00875, *Anisl* 7, and *Anisl* 22) have shown a genotype including the most common alleles found at those loci in the two parental species. The peaks representing the alleles found at those 100% diagnostic SSR loci in the parental taxa and in the three heterozygote specimens are reported in Fig. 3. Additionally, because the three larval specimens have shown a heterozygote genotype, also at the polymorphic sex-linked loci *Anisl* 4, *Anisl* 7 and *Anisl* 00134, they were assumed to be female worms.

According to the Bayesian clustering algorithm performed by STRUCTURE 2.3.3 (Fig. 4A), inferred from their genotypes recorded at those 100% diagnostic (i.e., *Anisl* 15, *Anisl* 2), and the partially diagnostic SSR loci (i.e., *Anisl* 10535, *Anisl* 05784 and *Anisl* 00875), plus those recorded at the *nas10* nDNA with the *K*= 2 clustering option, implemented by the Evanno

method, which best fitted the data set ($\Delta K = 12.80$) (Suppl. Fig.), a total of $N = 208$ specimens and $N = 133$ were assigned to the parental clusters (*A. pegreffii* and *A. berlandi* respectively) with high probability scores (Q-value = 100%). Of course, in this STRUCTURE analysis the other partially diagnostic sex-linked loci (i.e., *Anisl* 7 and *Anisl* 22) were excluded because of the hemizyosity of the male specimens. The three female nematodes have shown sign of mixed ancestry between the two groups having, in all cases, a Q-value = 0.50 (Fig. 4A). Finally, the analysis with NEWHYBRIDS identified with high level of confidence most of the individuals as 'pure' parental individuals of *A. pegreffii* or *A. berlandi* (Fig. 4B); the only exceptions were just those 3 individuals identified as admixed by the previous STRUCTURE analysis; the analysis assigned all these three specimens to the F1 hybrid class, with a 100% of probability value.

3.5. Intraspecific genetic differentiation of *A. pegreffii* and *A. berlandi*

Excluding the sex-linked loci because of the hemizyosity of males, the AMOVA analysis based on seven SSR markers showed that in both the species most of the variance was significantly allocated within individuals ($\approx 96\%$ and $\approx 97\%$). Further, a notable variation was found among individuals within populations, with $F_{IT} = 0.04$ ($p = 0.0009$) and $F_{IT} = 0.03$ ($p = 0.27$) in *A. pegreffii* and *A. berlandi*, respectively. (Table 6). Based on those SSR loci considered, a significantly high level of genetic differentiation has been found at the interspecific level, between the populations of *A. pegreffii* and those of *A. berlandi* (on average, $F_{st} \approx 0.605$, $p < 0.001$). In contrast, at the intraspecific level, lower and statistically not significant values were observed between metapopulations of larvae collected from fish species versus adult metapopulations of the two parasite species, recovered from cetacean hosts, being, $F_{st} \approx 0.002$ ($p = 0.21$) in *A. pegreffii*, and $F_{st} \approx 0.003$ ($p = 0.22$) in *A. berlandi*.

Analogously, considering the sequences analysis of the mitochondrial gene locus mtDNA *cox2*, the genetic differentiation among parental populations of *A. pegreffii* larvae and those adult

conspecifics was, on average, $F_{st} \approx 0.01$ ($p = 0.10$). A similar value, i.e., $F_{st} \approx 0.02$ ($p = 0.12$), was observed at the mitochondrial level, between parental populations of *A. berlandi* larvae with respect to adult samples analysed in the present study. Conversely, higher statistically significant pairwise comparison value at the mtDNA *cox2* sequences was recorded at the interspecific level between *A. pegreffii* and *A. berlandi* (i.e., on average, $F_{st} \approx 0.66$, $p < 0.001$).

4. Discussion

4.1 Nuclear markers to recognize parental *A. pegreffii* and *A. berlandi*, and their hybrid categories

In the present study, the testing of new SSR loci in species definition between *A. pegreffii* and *A. berlandi*. has disclosed two further nuclear loci (i.e., *Anisl 15* and *Anisl 2*) that were 100% diagnostic between the two species. These loci can identify both male and female worms of the two parasite taxa, at any developmental stage. The finding of these two loci having a high diagnostic power acquires particular importance in the discrimination analysis of the two parasite species, especially in sympatric areas. Indeed, among the SSR loci we had previously tested in Mattiucci et al., (2019), only one, i.e., the locus *Anisl 7*, was indicated as able to discriminate *A. berlandi* from *A. pegreffii* (Bello et al., 2020). However, since *Anisl 7* is a sex-linked locus, it cannot be included in a multigenotyping approach inferred by STRUCTURE analysis, to identify clusters of the two parental taxa; furthermore, it is not useful in recognising the potential occurrence of "male hybrid genotypes" between the two parental taxa. Conversely, the newly assessed loci *Anisl 15* and *Anisl 2*, were found to be fixed for two alternative alleles in both two species. These two loci were thus successfully used in the assignment of 208 *A. pegreffii* and 133 *A. berlandi* specimens with a 100% score value to their specific parental clusters, as well the 3 heterozygote specimens to their F1 hybrid category. The usefulness of the loci *Anisl 15* and *Anisl 2*, here investigated for the first time, were validated over a large number of specimens. They

were amongst those previously developed by Mladineo et al. (2017). These authors, however, did not recognise the discrimination power of those markers between the species of the *A. simplex* (s. l.) complex, nor their validity in disclosing hybrid categories and, even less, the presence of sex-linked loci. Indeed, in the last concerns, sex-linked loci SSR loci could be often erroneously considered as having null alleles, which is one among the reasons of significant departure from the HWE expectations when analysing the genetic variation inferred from DNA microsatellite loci (Criscione, et al., 2007a; Mattiucci et al., 2019).

The present genetic analysis has once again underlined the importance to use a multi-genotyping approach in the discrimination of parental taxa and admixed genotypes in a sympatric area of closely related species of *Anisakis*. Furthermore, this study has permitted a further validation of the nuclear gene locus *nas10* nDNA diagnostic power over a large number of larval specimens of the two parasite species. Indeed, in our previous study, the development of this nuclear marker was only investigated in adult specimens of *A. pegreffii* and *A. berlandi*, using the PCR-ARMS methodology based on the diagnostic positions detected as the *nas10* nDNA gene in the three species of the *A. simplex* (s. l.) complex (Palomba et al., 2020b). Here, the 344 specimens of *Anisakis* genotyped at the *nas10* nDNA, included in the Bayesian approach, produced complete agreement with the results obtained by the microsatellite loci. This congruency of these genetic markers permitted to define the population structure, in the contact zone, of the two parasite species, showing the existence of two distinct genetic clusters formed by the two parental species (i.e., *A. pegreffii* and *A. berlandi*), and providing clear evidence for their admixed genotypes (F1 hybrids *A. pegreffii* x *A. berlandi*) (Fig. 4 A, B). Thus, the present study represents the first demonstration of the occurrence of hybridization events of *A. pegreffii* and *A. berlandi* in a contact zone. Indeed, while interbreed events have been frequently reported in sympatric populations of *A. simplex* (s. s.) and *A. pegreffii* (Abollo et al., 2003; Marques et al., 2006; Umehara et al., 2006; Lee et al., 2009; Suzuki et al., 2010; Chou et al.,

2011; Molina-Fernández et al., 2015; Cipriani et al., 2015; Mattiucci et al., 2016; Gomez-Mateos et al., 2020), no hybridization or introgression phenomena had been so far reported between *A. simplex* (s. s.) and *A. berlandi* in contact and overlapping areas of their range of distribution (Mattiucci and Nascetti, 2007; Baldwin et al., 2011).

However, the finding of three F1 specimens of *A. pegreffii* and *A. berlandi* does not mean that the two taxa do not represent two "true biological species". In fact, the presence of only F1 hybrid genotypes does not imply that gene flow occurs between the two taxa, or that an incomplete speciation event had happened. It can instead indicate that reproductive isolation mechanisms may not be completely efficient for such related species. Pre-zygotic barriers for parasites include ecological isolation, isolation by host-specificity, as well as mate recognition systems (Southgate et al., 1998). In this regard, it seems that the first two conditions are not valid for *A. pegreffii* and *A. berlandi*, sharing both geographic and definitive hosts range (Mattiucci et al., 2018a). Thus, it might be supposed that chemical cues and/or other segregative mating recognition systems between conspecific adult specimens might not be highly efficient when syntopic infection in the same definitive host occurs. It can be also supposed that the absence of conspecific pairing partners, and mating stimuli for females of the rarer species in the syntopic situation may be important factors in increasing the likelihood of interspecific current hybridization (Avisé and Saunders, 1984). Thus, a tendency for hybridization would take place preferentially between parental species of parasites differing greatly in abundance. This could be the case for anisakid nematodes infecting the same definitive hosts in a contact zone. Indeed, taking into consideration that the number of mature females of anisakid parasites is generally higher than that of mature males in their definitive hosts (Ugland et al., 2004), when the intensity of the infection by one of these two species is low, mating events between the two species could occur, resulting in a first generation (F1) of larval hybrids (Mattiucci et al., 2016). In support of this hypothesis, considering the cetacean hosts here analysed, the occurrence of adult worms of

A. berlandi resulted lower with respect to those of the species *A. pegreffii* (Table 2). In addition, the F1 hybrids identified here have shown an *A. berlandi* maternal inheritance at the mtDNA *cox2* gene sequences analysis.

The frequency of current hybridization events between *A. pegreffii* and *A. berlandi* so far observed in this contact zone of the Pacific Ocean appears to be around 1%. Higher frequency (around 1,6%) of contemporary hybridization has been reported in the sympatric area of the Atlantic Iberian coast between the species *A. pegreffii* and *A. simplex* (s. s.) (Cipriani et al., 2015; Mattiucci et al., 2016). However, according to available data, the percentage of F1 adult worms between species of the *A. simplex* (s. l.) complex is far lower with respect to the number of F1 larval stage observed; indeed, only few hybrid adult worms were detected (Umehara et al., 2006; Cavallero et al., 2014). In those studies, a total of 5 hybrid mature female worms were found. The two larval specimens with F1 hybrid patterns here detected were female nematodes (i.e., showing heterozygote genotypes at the sex-linked loci), co-infecting the same individual fish host with parental specimens of both *A. pegreffii* and *A. berlandi*. This would suggest that some selective factors would decrease the F1 hybrid fitness, impeding the full development to mature adults worms in the definitive hosts of those larval F1 specimens, at the same rate of the parental species. Hybridization may give rise to F1 hybrids with low fertility or progeny's viability, preventing gene flow between parental taxa. The results so far obtained seem to reject the scenario of viable, i.e., fertile and able to reproduce F1 hybrids between the two taxa, because no backcross genotypes were identified in this study. Thus, the hybridization between *A. pegreffii* and *A. berlandi* could be evaluated as a sporadic and contemporary phenomenon.

Other examples of contemporary hybridization have been detected between *Ascaris lumbricoides* and *A. suum*, just discovered by a Bayesian clustering approach based on microsatellite data in sympatric populations from Guatemala and China (Criscione et al.,

2007b; Detwiler and Criscione, 2010), as well as in other cryptic parasite species, such as *Paramacrostrongylus* spp. (Chilton et al., 1997), *Fasciola hepatica* and *F. gigantica* (Agatsuma et al., 2000; Lin et al., 2007; Le et al., 2008; Peng et al., 2009; Calvani and Slapeta, 2020), *Schistosoma mansoni* and *S. rodhaini* (Morgan et al., 2003; Steinauer et al., 2010).

A study of such hybridization phenomena in parasites would shed light on epidemiological, ecological and phenotypic advantages eventually acquired by hybrid specimens. Hybridization events in a parasite species could have an impact on its adaptive radiation and diversification. Among these, a differential potential transmission of hybrids to a new host, the different adaptation to an intermediate host, or the colonization of a wider geographical range. These advantages could benefit hybrid genotypes with respect to the parental taxa, resulting in a possible adaptation to new or variation of biotic and abiotic factors (King et al., 2015).

The number of hybrid specimens so far detected between sibling species of the *A. simplex* (s. l.) complex species is too limited to make plausible assumptions about underlying ecological aspects. A more complete dataset could help to further elucidate these dynamics, by collecting and analysing the two parasite species in other hosts and from different basins where they geographically overlap.

The mtDNA *cox2* gene locus has proved its utility in the identification of all the species so far included in the genus *Anisakis* (Valentini et al., 2006), and in the phylogenetic constructions in concatenated analysis based on this locus (Mattiucci et al., 2018b). In addition, this molecular marker can add insight at the intraspecific level in the species of the *A. simplex* (s. l.) complex, concerning knowledge on possible population structuring and phylogeography (Baldwin et al., 2011; Mattiucci et al., 2018b), as well as in the estimating the genetic variability (Mattiucci et al., 2018a). However, its use as single marker in the identification of specimens of the species of the *A. simplex* (s. l.) complex, especially from sympatric areas, may be misleading in the discrimination of specimens between parental taxa and hybrid categories, also because of the

lack of information, so far, of recombination in the mitochondrial genome of these parasites. For instance, the detection of those three F1 hybrid genotypes would have been misidentified as *A. berlandi* if mtDNA *cox2* locus would have been the only locus tested in the present work (Table 2).

4.2 SSR sex-linked loci and their utility in gender determination of larval specimens

Sex-linked microsatellite loci are genes located on the X-chromosome. In the present work, four sex-linked loci (i.e., *Anisl 00314*, *Anisl 7*, *Anisl 4* and *Anisl 22*) have been discovered on the total panel of eleven DNA microsatellite markers so far studied in *A. pegreffii* and *A. berlandi*. As stated in the previous paragraph, the loci *Anisl 7* and *Anisl 22* are partially diagnostic between the two species. Therefore, when male and female specimens are available, it is advisable to verify if sex-linkage occurs in the SSR loci. To isolate a putative sex-linked locus, it is necessary to separate, on the basis of sex, the genotypes of the nematode sample. Where dealing with a sex-linked SSR locus, not all the genotypes observed can be likely used in the Bayesian clustering genotyping inference by STRUCTURE, irrespective of the sex identification. In fact, if the male specimens are homozygous at a considered microsatellite locus, it means that they are not “real” homozygous, but rather hemizygous. Having the chromosome(s) inherited from the mother only, without the chromosomal paternal counterpart, they possess just one copy of that gene. Therefore, taking into account that males are hemizygous (that is monoallelic), while females are heterozygous (that is biallelic), allele frequencies and allele frequencies-based analyses can be then calculated, as it has been carried out in the present study.

While the proportion of sex-linked microsatellites on the whole genome of the species of *A. simplex* (s. l.) was roughly estimated around the 25% in our previous SSRs analysis (Mattiucci et al., 2019), the present study updates this percentage, that should be considered up to 33%, due to the scoring of further DNA microsatellite loci. This value represents a higher percentage in

comparison to the 20% that Criscione et al. (2007a) supposed for ascarids (such as *Ascaris lumbricoides*). To date, karyotypes of anisakid nematodes were not determined. In ascaridoid nematodes, males are XO, while females are XX. Maternal gametes always contain an X chromosome, so the sex of the "offspring" specimens depends on whether a sex chromosome occurs in the male gamete. Its sperm normally contains either X chromosome or no sex chromosome at all. In the evolution of sex chromosomes of parasitic nematodes, it is retained that the XO sex chromosome system is the ancestral state in nematodes, while the XY is the derived state (Foster et al., 2020). Some nematodes such as the non-parasitic *Caenorhabditis elegans* have a XO sex-determination system, while others, such as the filarial parasite *Brugia malayi*, have an XY mechanism (Foster et al., 2020). However, the only reference available for ascaridoid nematodes regards *Ascaris suum*, which shows 19 autosomes and 5X chromosomes ($2n = 38A + 10X$ in females; $38A + 5X$ in males) (Müller and Tobler, 2000). Future cytological work would shed light on the sex chromosomes of anisakid parasites and their evolutionary significance.

Besides being important for population genetics studies, microsatellites can be also a useful and powerful tool to distinguish females from males in organisms with XO sex determination (Harvey and Viney, 2001). The sex-associated SSR markers so far disclosed in the species of *A. simplex* (s. l.) complex and in the present study, can be of help in revealing an alternative mode of sex identification in their developmental stages (Mattiucci et al., 2019). In fact, if male hemizyosity is assumed, sex-linked loci could likely allow identification of any female at any life-history stage (even as L3 larva or L4 immature worm), when a larval individual shows a heterozygote genotype at one or more of those loci. According to this assumption, for instance, the heterozygote genotypes found, at least, in one of the four sex-linked loci, would allow the identification as female N= 55 of L3 and L4 stages of *A. pegreffii* analysed. Similarly, N= 52 of L3 and L4 larval stages of *A. berlandi*, being heterozygotes at the loci *Anisl 7* and/or *Anisl 4* (the

other two loci *Anisl* 00314 and *Anisl* 22 are monomorphic in this species), can be assumed to be female. However, the occurrence of homozygote genotypes in females could also occur. Thus, in future studies, estimating the statistical occurrence of homozygous genotypes in large number of adult females on as many as possible sex-linked loci, a high probability of correct sex identification of third and fourth larval stage specimens of these parasite species could be reached.

4.3 Genetic differentiation and variability of *A. pegreffii* and *A. berlandi*

The genetic/molecular results obtained further confirmed that the two studied taxa are genetically distinct in both nuclear and mitochondrial markers. The multilocus genotyping approach (STRUCTURE) based on nuclear genes of the two parasite taxa has demonstrated that the analysed metapopulations, despite the wide host range, were strongly associated with two distinct panmictic units, corresponding, respectively, to the two species *A. pegreffii* and *A. berlandi*. The result was also corroborated by the pairwise F_{ST} estimates found at the interspecific level, at both nuclear (as based on SSR loci) and mitochondrial (mtDNA *cox2*) gene level.

Whereas, a high genetic homogeneity across sample populations of the two species collected from different fish and cetacean host species was observed in both *A. pegreffii* and *A. berlandi* occurring in the contact zone of New Zealand waters. The observed values are in agreement with those previously recorded, among populations of the same species, in samples from other geographic areas, also based on DNA microsatellite loci (Mattiucci et al., 2019) and mtDNA *cox2* sequences analysis (Mattiucci et al., 2014).

At the intrapopulation level, previous scoring of SSR loci based on adult populations of *A. berlandi* have detected F_{IT} values higher than zero (Bello et al., 2020), likely suggesting a possible certain subdivision between subpopulations of this parasite species. Whereas, in the

present study, the additional scoring of a larger number of specimens, including larval forms from several fish species, has shown a not significant F_{IT} , and F_{IS} values, being those very close to zero (Table 6). This result provides no evidence for deviation from random mating within intrapopulations of *A. berlandi* from this area of the Pacific, i.e., the existing of panmixia. The partitioning of genetic differentiation between and within the populations of *A. pegreffii* studied at the SSRs loci, have shown that most of the diversity is within populations, being also the F_{ST} among populations very close to zero (Table 6). This again implies the existence of a high level of gene flow between the populations of *A. pegreffii* inhabiting New Zealand waters, despite the high genetic polymorphism presently disclosed at 11 DNA microsatellite loci. At the intraspecific level, a moderate genetic differentiation (on average, $F_{ST}= 0.060$) estimated on 7 SSR loci, has been posited between the Boreal populations of *A. pegreffii* from the Mediterranean Sea, and their conspecifics from the New Zealand southern Pacific and Argentine Southern Atlantic (Mattiucci et al., 2019). The parasite population genetic substructuring would be also driven by the population genetic structure of its definitive hosts (Esch and Fernandez, 1993). In the present case, the population structure of some cetacean species, from different geographical areas would also be responsible for maintaining such level of genetic substructuring of their endoparasites (i.e., *A. pegreffii*) in Boreal (i.e., Mediterranean Sea) and Austral (i.e., New Zealand) waters. Indeed, for instance, the genetic diversity and population genetic structure of the bottlenose dolphin *Tursiops truncatus* identify its New Zealand population as the most divergent from populations of the Atlantic and Mediterranean waters. Despite this, *T. truncatus* maintains high level of gene flow over long distances (Tezanos-Pinto et al., 2009). This finding seems to mirror the population genetic substructuring so far detected in *A. pegreffii* from different geographical areas included in its range of distribution, as the possible effect of host-parasite relationship. The possible role of biotic factors, in terms of host population density, vagility, and genetic population structure, has been

indeed suggested to be a driver in the epidemiology and the genetic structure of the parasites species included in the *A. simplex* (s. l.) complex (Mattiucci et al., 2018a).

Despite the presence of larval stages of *A. berlandi* from fish host has been recorded in several areas of the Southern region, and recently also from the Argentine coast (Irigoitia et al., 2018), sufficient data on its population genetic structure investigated by DNA microsatellite markers and mtDNA *cox2* are not yet available. This prevents meaningful discussion about possible spatial genetic diversity between different southern populations of *A. berlandi*. Future analysis of nuclear and mitochondrial gene distributions in populations of the two parasite species from different Boreal and Southern Ocean waters will help provide a worldwide perspective on their population structure and genetic diversity.

Multi-host ascaridoid populations are expected to maintain higher levels of genetic polymorphism than parasites with a narrow host range, because they are unlikely to experience extreme population number fluctuations (Bullini et al., 1998; Mattiucci and Nascetti, 2007, 2008). The results obtained in the present study agree with this prediction, as the polymorphism detected was high at all microsatellite loci, in the scored populations of *A. pegreffii* (on average, $He = 0.56$); a lower value was observed in *A. berlandi* (on average, $He = 0.28$). A high genetic polymorphism was previously found at other nuclear loci (allozymes) in the populations of the two parasite species from New Zealand waters, with average values of $He = 0.11$ in *A. pegreffii*, and $He = 0.24$ in *A. berlandi* (Mattiucci et al., 1997). High genetic variability has been generally reported in species of the *A. simplex* (s. l.) complex, also inferred from other nuclear gene loci, such as allozymes (Mattiucci and Nascetti, 2008). This finding could be ascribed to the large effective population size of anisakid species generally observed in both intermediate/paratenic (fish) and definitive (cetaceans) hosts. Furthermore, a high level of gene flow exists between conspecific populations of these parasites, maintained by the high vagility of many of their hosts, carrying the parasites over large migration routes, across distant oceanic basins. In the present

study, parasitic infection levels exhibited by the two parasite species in different fish species are not given. Additional genetic studies and data on the parasite population density of *A. pegreffii* and *A. berlandi* from different geographic areas of the Southern Region are required, in order to investigate spatial genetic structure of the two parasite species, their larval and adult abundance, their effective population sizes and genetic variability.

5. CONCLUDING REMARKS

This is the first study employing a wide-genotyping approach to identify adults and larvae of two sibling species of the *A. simplex* (s. l.) complex originating from New Zealand waters. Several *A. simplex* (s. l.) nuclear gene loci, discovered recently and also in this work, have a clear diagnostic capacity to identify *A. pegreffii* and *A. berlandi*. Markers based on these loci have allowed genetic data collection, which elaborated by updated robust Bayesian methodologies, have dramatically raised the possibility of distinguishing parental and hybrid genotypes, and genetically characterise populations of these nematodes. Indeed, despite the fact that DNA microsatellite loci are considered as markers under selection, the scored SSR loci could be used as markers of neutral genetic structure of the two parasite species. All the tested loci have indeed given a similar signal and behave similarly in the analysis. Thus, it appears that these SSR loci reflect the genetic structure of the regions in the genome of those parasite species that are not likely involved in the host–parasite interaction, i.e., regions of the genome that are not under selection.

It has become clear that without a wider investigation of the genetic architecture of these anisakid parasites, based on several nuclear genes, misinterpretation of naturally occurring hybridization and/or introgression events can occur. Indeed, hybridization does not necessarily generate permanent genetic exchange between the taxa; it could be a transient phenomenon. When, on the contrary, interspecific mating events occur between hybrids and their parental

species or between hybrid individuals, both events resulting in backcross individuals, a more permanent introgression phenomenon would exist, which could also represent an adaptive trait from one species into the other. However, the current ambiguity surrounding the distinction between hybridization and introgression in the species of *A. simplex* (s. l.) complex, such as for instance, between *A. simplex* (s. s.) and *A. pegreffii*, is mainly due to a tendency to investigate a single locus when, instead, a multi-loci or whole-genome approach would be required.

In the present study, diagnostic genotyping at other nuclear markers, other than those inferred from SSR loci, such as those gathered from metallopeptidase *nas10* nDNA, will help clarify the affinities and genetic structure of anisakid parasites and their populations in sympatric areas. The combined nuclear data sets have allowed the successful detection of current hybridisation between *A. pegreffii* x *A. berlandi*. However, investigations at other nuclear loci, even starting from an available draft genome of *A. simplex* (s. l.) (Łopieńska-Biernat et al., 2019), should be pursued to disclose further diagnostic SNPs at certain gene loci, to aid in differentiating and mapping the hybridization/introgression events between the three species of the *A. simplex* (s. l.) complex in their contact areas. This will also add insights into the phenotypic and functional implications of naturally occurring hybrids between the three taxa. In addition, future studies employing as many as possible sex-linked loci would be also aimed for sex identification of third and fourth larval stage specimens of these worms. Sex determination in these parasites will facilitate further investigations, such as the significance of relative proportions of adult males/females in definitive hosts in comparison with larval stages occurring in intermediate/paratenic hosts, differential sites of infection in fish hosts, and the effect of differential gene expression levels between male and female specimens. In the last concerns, for instance, differential gene transcriptions between L3 and L4 in *A. pegreffii* and *A. simplex* (s. s.) (Kim et al., 2018) has been demonstrated, as well as the existence of differential gene expression of adaptive genes in those parasite species infecting fish host tissues (Palomba et al., 2020a).

In conclusion, the new and emerging wide-genotyping approach in the species of the *A. simplex* (s. l) complex could provide information also in their phenotypic and ecological traits, which, in turn, would have also epidemiological and pathological implications. These last aspects acquire human health importance because, while currently *A. pegreffii* and *A. simplex* (s. s.) are known as the etiological agents of human anisakiasis, the zoonotic role of *A. berlandi* is so far unknown (Mattiucci et al., 2018a).

Authors contribution

Conceptualization, SM, GN, EB, MaP; methodology, EB, MaP, MiP; material sampling: SCW; writing-original draft preparation, SM, EB, MaP, PC, SCW; writing-review and editing, SM, PC SCW; supervision, SM, GN. All authors have read and agreed to the published version of the manuscript.

Acknowledgements

We thank Prof. Wendi Roe (Massey University), Manawhenua Ki Mohua, Mike Ogle (Department of Conservation), Dr Felix Marx (Otago Museum) and Julie Nicholls (Moana, NZ) for providing specimens. In addition, the Authors are grateful to Dr Armando Macali from the Department of Biological and Ecological Sciences (Tuscia University, Italy) for sampling larval nematodes from fish species included in this study.

Funding

The work was partially supported by a grant from MIUR-PNRA2016_00279. Additionally, S Mattiucci carried out part of this research work with a grant from "Istituto Pasteur Italia-Fondazione Cenci Bolognetti", Italy.

Conflicts of Interest

The authors declare no conflict of interest.

Legends to the figures

Fig 1. Alignment of the *nas10* nDNA gene locus for the two species *A. pegreffii* and *A. berlandi*.

The arrows show the fixed diagnostic nucleotide position detected between the two species. The heterozygote pattern at the diagnostic nucleotide position is shown. Dots indicate identity; standard IUPAC codes were used, i.e., S = C/G

Fig 2. (A) F_{IS} value calculated at nine microsatellite loci (except *Anisl 15* and *Anisl 2*, because of the monomorphism in both the species) studied in the two species *A. pegreffii* and *A. berlandi*.

(B) F_{IS} in male and female adult specimens at the sex-linked loci in *A. pegreffii* (i.e., *Anisl 00314*, *Anisl 7*, *Anisl 4* and *Anisl 22*) and in *A. berlandi* (i.e., *Anisl 7* and *Anisl 4*). Negative values indicate heterozygous excess, while positive values indicate homozygous excess from that expected under Hardy–Weinberg Equilibrium (HWE).

Fig. 3. Electropherograms and observed genotypes of pure parental *A. berlandi* (on the left), F1 hybrids (in the middle) and pure parental *A. pegreffii* (on the right) at the diagnostic loci (100%) between the two *Anisakis* species (i.e., *Anisl 15* and *Anisl 2*); Discriminant alleles for *A. berlandi* are given as empty peaks; discriminant alleles for *A. pegreffii* are shown as full peaks.

Fig. 4. Percentage contribution (Q value) of *A. pegreffii* and *A. berlandi* species to the multilocus genotype of each studied individual (barplot), estimated at five SSR loci (i.e., *Anisl 10535*, *Anisl 05784*, *Anisl 00875*, *Anisl 15* and *Anisl 2*) and at the nuclear gene *nas10*, by using STRUCTURE

with $k = 2$ (A), and probability of assignment to a particular hybrid class estimated using NEWHYBRIDS (B) (Anderson and Thompson, 2002).

Suppl. Fig. Estimate of the best number of clusters (K) describing the population structure of the *Anisakis* spp. samples analyzed in the present study. The computation was carried out by using STRUCTURE HARVESTER. (A) highest ln-probability and (B) DeltaK optimality criteria.

References

- Abollo, E., Paggi, L., Pascual, S., D'Amelio, S., 2003. Occurrence of recombinant genotypes of *Anisakis simplex* s.s. and *Anisakis pegreffii* (Nematoda: Anisakidae) in an area of sympatry. Infect. Genet. Evol. 3, 175-181. [https://doi.org/10.1016/S1567-1348\(03\)00073-X](https://doi.org/10.1016/S1567-1348(03)00073-X).
- Agatsuma, T., Arakawa, Y., Iwagami, M., Honzako, Y., Cahyaningsih, U., Kang, S., Hong, S., 2000. Molecular evidence of natural hybridization between *Fasciola hepatica* and *F. gigantica*. Parasitol. Int. 49 (3), 231-238. [https://doi.org/10.1016/S1383-5769\(00\)00051-9](https://doi.org/10.1016/S1383-5769(00)00051-9).
- Anderson, E.C., Thompson, E.A., 2002. A model-based method for identifying species hybrids using multilocus genetic data. Genetics. 160 (3), 1217-1229.
- Awise, J.C., Saunders, N.C., 1984. Hybridization and introgression among species of sunfish (*Lepomis*): analysis by mitochondrial DNA and allozyme markers. Genetics 108, 237-255. <https://doi.org/10.1093/genetics/108.1.237>.
- Baldwin, R.E., Rew, M.B., Johansson, M.L., Banks, M.A., Jacobson, K.C., 2011. Population structure of three species of *Anisakis* nematodes recovered from Pacific sardines (*Sardinops*

sagax) distributed throughout the California Current System. J. Parasitol. 97, 545–554.
<https://doi.org/10.1645/GE-2690.1>.

Bello, E., Paoletti, M., Webb, S.C., Nascetti, G., Mattiucci, S., 2020. Cross-species utility of
microsatellite loci for the genetic characterisation of *Anisakis berlandi* (Nematoda:
Anisakidae). Parasite 27, 9. <https://doi.org/10.1051/parasite/2020004>.

Berland, B., 1961. Nematodes from some Norwegian marine fishes. Sarsia 2, 1-50.
<https://doi.org/10.1080/00364827.1961.10410245>.

Bullini, L., Nascetti, G., Paggi, L., Orecchia, P., Mattiucci, S., Berland, B., 1986. Genetic variation
of ascaridoid worms with different life cycles. Evolution, 40(2), 437-440.

Cavallero, S., Nadler, S.A., Paggi, L., Barros, N.B., D'Amelio, S., 2011. Molecular characterization
and phylogeny of anisakid nematodes from cetaceans from southeastern Atlantic coasts of
USA, Gulf of Mexico and Caribbean Sea. Parasitol. Res. 108, 781–792.
<https://doi.org/10.1007/s00436-010-2226-y>.

Cavallero, S., Costa, A., Caracappa, S., Gambetta, B., D'Amelio, S., 2014. Putative hybrids
between two *Anisakis* cryptic species: Molecular genotyping using High Resolution Melting.
Exp. Parasitol. 146, 87-93. <https://doi.org/10.1016/j.exppara.2014.08.017>.

Calvani, N.E.D., Šlapeta, J., 2020. *Fasciola* Species Introgression: Just a Fluke or Something
More? Trends Parasitol. 37, 25.34. <https://doi.org/10.1016/j.pt.2020.09.008>.

Cipriani, P., Smaldone, G., Acerra, V., D'Angelo, L., Anastasio, A., Bellisario, B., Palma, G.,
Nascetti, G., Mattiucci, S., 2015. Genetic identification and distribution of the parasitic larvae
of *Anisakis pegreffii* and *Anisakis simplex* (s. s.) in European hake *Merluccius merluccius* from

the Tyrrhenian Sea and Spanish Atlantic coast: implications for food safety. Int. J. Food Microbiol. 198, 1-8. <https://doi.org/10.1016/j.ijfoodmicro.2014.11.019>.

Chilton, N.B., Beveridge, I., Hoste, H., Gasser, R.B., 1997. Evidence for hybridisation between *Paramacropostrongylus iugalis* and *P. typicus* (Nematoda: Strongyloidea) in grey kangaroos, *Macropus fuliginosus* and *M. giganteus*, in a zone of sympatry in eastern Australia. Int. J. Parasitol. 27, 475-482. [https://doi.org/10.1016/S0020-7519\(97\)00024-6](https://doi.org/10.1016/S0020-7519(97)00024-6).

Chou, Y.Y., Wang, C.S., Chen, H.G., Chen, H.Y., Chen, S.N., Shih, H.H., 2011. Parasitism between *Anisakis simplex* (Nematoda: Anisakidae) third-stage larvae and the spotted mackerel *Scomber australasicus* with regard to the application of stock identification. Vet. Parasitol. 177, 324-331. <https://doi.org/10.1016/j.vetpar.2010.12.003>.

Criscione, C.D., Anderson, J.D., Raby, K., Sudimack, D., Subedi, J., Rai, D.R., Upadhayay, R.P., Jha, B., Williams-Blangero, S., Anderson, T.J., 2007a. Microsatellite markers for the human nematode parasite *Ascaris lumbricoides*: development and assessment of utility. J. Parasitol. 93, 704-708. <https://doi.org/10.1645/GE-1058R.1>.

Criscione, C.D., Anderson, J.D., Sudimack, D., Peng, W., Jha, B., Williams-Blangero, S., Anderson, T.J., 2007b. Disentangling hybridization and host colonization in parasitic roundworms of humans and pigs. Proc. Biol. Sci. 274 (1626), 2669-2677. <https://doi.org/10.1098/rspb.2007.0877>.

D'Amelio, S., Mathiopoulos, K.D., Santos, C.P., Pugachev, O.N., Webb, S.C., Picanço, M., Paggi, L., 2000. Genetic markers in ribosomal DNA for the identification of members of the genus *Anisakis* (Nematoda: Ascaridoidea) defined by polymerase chain reaction-based restriction fragment length polymorphism. Int. J. Parasitol. 30, 223-226. [https://doi.org/10.1016/S0020-7519\(99\)00178-2](https://doi.org/10.1016/S0020-7519(99)00178-2).

- Detwiler, J.T., Criscione, C.D., 2010. An Infectious Topic in Reticulate Evolution: Introgression and Hybridization in Animal Parasites. *Genes* 1, 102-123. <https://doi.org/10.3390/genes1010102>.
- Esch, G.W., Fernández, J.C. (1993) Factors influencing parasite populations, in: Esch, G.W., Fernández, J.C. (Eds.), *A Functional Biology of Parasitism*. Functional Biology Series. Springer, Dordrecht. https://doi.org/10.1007/978-94-011-2352-5_3
- Evanno, G., Regnaut, S., Goudet, J., 2005. Detecting the number of clusters of individuals using the software STRUCTURE: a simulation study. *Mol. Ecol.* 14, 2611–2620. <https://doi.org/10.1111/j.1365-294X.2005.02553.x>.
- Excoffier, L., Lischer, H.E., 2010. Arlequin suite ver. 3.5: a new series of programs to perform population genetics analyses under Linux and Windows. *Mol. Ecol. Resour.* 10, 564-567. <https://doi.org/10.1111/j.1755-0998.2010.02847.x>.
- Foster, J.M, Grote, A., Mattick, J., Tracey, A., Tsai, Y., Chung, M., Cotton, J.A., Clark, T.A., Geber, A., Holroyd, N., Korlach, J., Li, Y., Libro, S., Lustigman, S., Michalski, M.L., Paulini, M., Rogers, M.B., Teigen, L., Twaddle, A., Welch, L., Berriman, M., Dunning Hotopp, J.C., Ghedin, E., 2020. Sex chromosome evolution in parasitic nematodes of humans. *Nat. Commun.* 11 (1), 1-12. <https://doi.org/10.1038/s41467-020-15654-6>.
- Gómez-Mateos, M., Merino-Espinosa, G., Corpas-López, V., Valero-López, A., Martín-Sánchez, J., 2020. A multi-restriction fragment length polymorphism genotyping approach including the beta-tubulin gene as a new differential nuclear marker for the recognition of the cryptic species *Anisakis simplex* s.s. and *Anisakis pegreffii* and their hybridization events. *Vet. Parasitol.* 283, 109162. <https://doi.org/10.1016/j.vetpar.2020.109162>.

- Gomes, T.L., Quiazon, K.M., Kotake, M., Fujise, Y., Ohizumi, H., Itoh, N., Yoshinaga, T., 2021. *Anisakis* spp. in toothed and baleen whales from Japanese waters with notes on their potential role as biological tags. *Parasitol. Int.* 80, 102228. <https://doi.org/10.1016/j.parint.2020.102228>.
- Gomes, T.L., Quiazon, K.M., Kotake, M., Itoh, N., Yoshinaga, T., 2020. *Anisakis* spp. in fishery products from Japanese waters: Updated insights on host prevalence and human infection risk factors. *Parasitol. Int.* 78, 102137. <https://doi.org/10.1016/j.parint.2020.102137>.
- Harvey, S.C., Viney, M.E., 2001. Sex determination in the parasitic nematode *Strongyloides ratti*. *Genetics*. 158 (4), 1527-1533.
- Irigoitia, M.M., Braicovich, P.E., Lanfranchi, A.L., Farber, M.D., Timi, J.T., 2018. Distribution of anisakid nematodes parasitizing rajiform skates under commercial exploitation in the Southwestern Atlantic. *Int. J. Food Microbiol.* 267, 20-28. <https://doi.org/10.1016/j.ijfoodmicro.2017.12.009>.
- Kim, J.H., Kim, J.O., Jeon, C.H., Nam, U.H., Subramaniyan, S., Yoo, S.I., Park, J.H., 2018. Comparative transcriptome analyses of the third and fourth stage larvae of *Anisakis simplex* (Nematoda: Anisakidae). *Mol. Biochem. Parasitol.* 226, 24-33. <https://doi.org/10.1016/j.molbiopara.2018.10.002>.
- King, K.C, Stelken, R.B., Webster, J.P., Smith, D.F., Brockhurst, M.A. (2015). Hybridization in parasites: Consequences for adaptive evolution, pathogenesis, and public health in a changing world. *PLOS Pathogens*, 3: 12- | DOI:10.1371/journal.ppat.1005098
- Klimpel, S., Busch, M.W., Kuhn, T., Rohde, A., Palm, H.W., 2010. The *Anisakis simplex* complex off the South Shetland Islands (Antarctica): endemic populations versus introduction through migratory hosts. *Mar. Ecol. Prog. Ser.* 403, 1–11. <https://doi.org/10.3354/meps08501>.

- Larkin, M.A., Blackshields, G., Brown, N.P., Chenna, R., McGettigan, P.A., McWilliam, H.,
Valentin, F., Wallace, I.M., Wilm, A., Lopez, R., Thompson, J.D., Gibson, T.J., Higgins, D.G.,
2007. Clustal W and Clustal X version 2.0. *Bioinformatics* 23, 2947–2948.
<https://doi.org/10.1093/bioinformatics/btm404>.
- Le, T.H., Van De, N., Agatsuma, T., Nguyen, T.G.T., Nguyen, Q.D., McManus, D.P., Blair, D.,
2008. Human fascioliasis and the presence of hybrid/introgressed forms of *Fasciola hepatica*
and *Fasciola gigantica* in Vietnam. *Int. J. Parasitol.* 38 (6), 725-730.
<https://doi.org/10.1016/j.ijpara.2007.10.003>.
- Lee, M.H., Cheon, D., Choi, C., 2009. Molecular genotyping of *Anisakis* species from Korean sea
fish by polymerase chain reaction–restriction fragment length polymorphism (PCR-RFLP).
Food Control 20, 623-626. <https://doi.org/10.1016/j.foodcont.2008.09.007>.
- Lin, R.Q., Dong, S.J., Nie, K., Wang, C.R., Song, H.Q., Li, A.X., Zhu, X.Q., 2007. Sequence
analysis of the first internal transcribed spacer of rDNA supports the existence of the
intermediate *Fasciola* between *F. hepatica* and *F. gigantica* in mainland China. *Parasitol. Res.*
101 (3), 813-817. <https://doi.org/10.1007/s00436-007-0512-0>.
- Łopieńska-Biernat, E., Paukszto, L., Jastrzebski, J.P., Myszczyński, K., Polak, I., Stryński, R.,
2019. Genome-wide analysis of *Anisakis simplex* sensu lato: The role of carbohydrate
metabolism genes in the parasite's development. *Int. J. Parasitol.* 49, 933-94.
<https://doi.org/10.1016/j.ijpara.2019.06.006>.
- Marques, J.F., Cabral, H.N., Busi, M., D'Amelio, S., 2006. Molecular identification of *Anisakis*
species from Pleuronectiformes off the Portuguese coast. *J. Helminthol.* 80 (1), 47-51.
<https://doi.org/10.1079/JOH2005325>.

- Mattiucci, S., Nascetti, G., Cianchi, R., Paggi, L., Arduino, P., Margolis, L., Bratney, J., Webb, S.C., D'Amelio, S., Orecchia, P., Bullini, L., 1997. Genetic and ecological data on the *Anisakis simplex* complex with evidence for a new species (Nematoda, Ascaridoidea, Anisakidae). J. Parasitol. 83 (3), 401-416. <https://doi.org/10.2307/3284402>.
- Mattiucci, S., Nascetti, G., 2007. Genetic diversity and infection levels of anisakid nematodes parasitic in fish and marine mammals from Boreal and Austral hemispheres. Vet. Parasitol. 148, 43-57. <https://doi.org/10.1016/j.vetpar.2007.05.009>.
- Mattiucci, S., Nascetti, G., 2008. Advances and trends in the molecular systematics of *Anisakis* nematodes, with implications for their evolutionary ecology and host-parasite co-evolutionary processes. Adv. Parasitol. 66, 47-148. [https://doi.org/10.1016/S0065-308X\(08\)00202-9](https://doi.org/10.1016/S0065-308X(08)00202-9).
- Mattiucci, S., Cipriani, P., Webb, S.C., Paoletti, M., Marcer, F., Bellisario, B., Gibson, D.I., Nascetti, G., 2014. Genetic and morphological approaches distinguishing the three sibling species of the *Anisakis simplex* species complex, with a species designation as *Anisakis berlandi* n. sp. for *A. simplex* sp. C (Nematoda: Anisakidae). J. Parasitol. 15, 12-15. <https://doi.org/10.1645/12-120.1>.
- Mattiucci, S., Acerra, V., Paoletti, M., Cipriani, P., Levsen, A., Webb, S.C., Canestrelli, D., Nascetti, G., 2016. No more time to stay 'single' in the detection of *Anisakis pegreffii*, *A. simplex* (s. s.) and hybridization events between them: a multi-marker nuclear genotyping approach. Parasitology 143, 998-1011. <https://doi.org/10.1017/S0031182016000330>.
- Mattiucci, S., Cipriani, P., Levsen, A., Paoletti, M., Nascetti, G., 2018a. Molecular epidemiology of *Anisakis* and anisakiasis: an ecological and evolutionary road map. Adv. Parasitol. 99, 93-263. <https://doi.org/10.1016/bs.apar.2017.12.001>.

- Mattiucci, S., Giulietti, L., Paoletti, M., Cipriani, P., Gay, M., Levsen, A., Paoletti, M., Nascetti, G., 2018b. Population genetic structure of the parasite *Anisakis simplex* (s. s.) collected in *Clupea harengus* L. from North East Atlantic fishing grounds. Fish. Res. 202, 103–111. <https://doi.org/10.1016/j.fishres.2017.08.002>.
- Mattiucci, S., Bello, E., Paoletti, M., Webb, S.C., Timi, J.T., Levsen, A., Cipriani, P., Nascetti, G., 2019. Novel polymorphic microsatellite loci in *Anisakis pegreffii* and *A. simplex* (s. s.) (Nematoda: Anisakidae): implications for species recognition and population genetic analysis. Parasitology 1–17. <https://doi.org/10.1017/S003118201900074X>.
- Mladineo, I., Trumbić, Ž., Radonić, I., Vrbatović, A., Hrabar, J., Bušelić, I., 2017. *Anisakis simplex* complex: ecological significance of recombinant genotypes in an allopatric area of the Adriatic Sea inferred by genome derived simple sequence repeats. Int. J. Parasitol. 47, 215–223. <https://doi.org/10.1016/j.ijpara.2016.11.003>.
- Mayr, E., 1963. Animal Species and Evolution. Harvard University Press, Cambridge.
- Molina-Fernández, D., Malagón, D., Gómez-Mateos, M., Benítez, R., Martín-Sánchez, J., Adroher, F.J., 2015. Fishing area and fish size as risk factors of *Anisakis* infection in sardines (*Sardina pilchardus*) from Iberian waters, south-western Europe. Int. J. Food Microbiol. 203, 27–34. <https://doi.org/10.1016/j.ijfoodmicro.2015.02.024>.
- Morgan, J.A., DeJong, R.J., Lwambo, N.J., Mungai, B.N., Mkoji, G.M., Loker, E.S., 2003. First report of a natural hybrid between *Schistosoma mansoni* and *S. rodhaini*. J. Parasitol. 89 (2), 416–419. [https://doi.org/10.1645/0022-3395\(2003\)089\[0416:FROANH\]2.0.CO;2](https://doi.org/10.1645/0022-3395(2003)089[0416:FROANH]2.0.CO;2).
- Müller, F., Tobler, H., 2000. Chromatin diminution in the parasitic nematodes *Ascaris suum* and *Parascaris univalens*. Int. J. Parasitol. 30, 391–399. [https://doi.org/10.1016/S0020-7519\(99\)00199-X](https://doi.org/10.1016/S0020-7519(99)00199-X).

- Nascetti, G., Paggi, L., Orecchia, P., Smith, J.W., Mattiucci, S., Bullini, L., 1986. Electrophoretic studies on *Anisakis simplex* complex (Ascaridida: Anisakidae) from the Mediterranean and North East Atlantic. *Int. J. Parasitol.* 16, 633–640. [https://doi.org/10.1016/0020-7519\(86\)90032-9](https://doi.org/10.1016/0020-7519(86)90032-9).
- Palomba, M., Cipriani, P., Giulietti, L., Levsen, A., Nascetti, G., Mattiucci, S., 2020a. Differences in Gene Expression Profiles of Seven Target Proteins in Third-Stage Larvae of *Anisakis simplex* (Sensu Stricto) by Sites of Infection in Blue Whiting (*Micromesistius poutassou*). *Genes* 11 (5), 559. <https://doi.org/10.3390/genes11050559>.
- Palomba, M., Paoletti, M., Webb, S.C., Nascetti, G., Mattiucci, S., 2020b. A novel nuclear marker and development of an ARMS-PCR assay targeting the metallopeptidase 10 (*nas* 10) locus to identify the species of the *Anisakis simplex* (s. l.) complex (Nematoda, Anisakidae). *Parasite* 27, 39. <https://doi.org/10.1051/parasite/2020033>.
- Peng, M., Ichinomiya, M., Ohtori, M., Ichikawa, M., Shibahara, T., Itagaki, T., 2009. Molecular characterization of *Fasciola hepatica*, *Fasciola gigantica*, and aspermic *Fasciola* sp. in China based on nuclear and mitochondrial DNA. *Parasitol. Res.* 105 (3), 809. <https://doi.org/10.1007/s00436-009-1459-0>.
- Pritchard, J.K., Stephens, M., Donnelly, P., 2000. Inference of population structure using multilocus genotype data. *Genetics* 155, 945-959.
- Shamsi, S., Gasser, R., Beveridge, I., 2012. Genetic characterisation and taxonomy of species of *Anisakis* (Nematoda: Anisakidae) parasitic in Australian marine mammals. *Invertebr. Syst.* 26, 204-212. <https://doi.org/10.1071/IS11019>.

- Shamsi, S., Briand, M.J., Justine, J.L., 2017. Occurrence of *Anisakis* (Nematoda: Anisakidae) larvae in unusual hosts in Southern hemisphere. *Parasit. Int.* 66 (6), 835-840. <https://doi.org/10.1016/j.parint.2017.08.002>
- Southgate, V.R., Jourdane, J., Tchuem, Tchente, L.A., 1998. Recent studies on the reproductive biology of the schistosomes and their relevance to speciation in the Digenea. *Int. J. Parasitol.* 28 (8), 1159-1172. [https://doi.org/10.1016/S0020-7519\(98\)00021-6](https://doi.org/10.1016/S0020-7519(98)00021-6).
- Steinauer, M.L., Blouin, M.S., Criscione, C.D., 2010. Applying evolutionary genetics to schistosome epidemiology. *Infect. Genet. Evol.* 10, 433-443. <https://doi.org/10.1016/j.meegid.2010.02.007>.
- Suzuki, J., Murata, R., Hosaka, M., Araki, J., 2010. Risk factors for human *Anisakis* infection and association between the geographic origins of *Scomber japonicus* and anisakid nematodes. *Int. J. Food Microbiol.* 137, 88-93. <https://doi.org/10.1016/j.ijfoodmicro.2009.10.001>.
- Tezanos-Pinto, G., Scott Baker, C., Russell, K., Martien, K., Baird, R.W., Hutt, A., Stone, G., Mignucci-Giannoni, A.A., Caballero, S., Endo, T., Lavery, S., Oremus, M., Olavarria, C., Garrigue, C., 2009. A Worldwide Perspective on the Population Structure and Genetic Diversity of Bottlenose Dolphins (*Tursiops truncatus*) in New Zealand. *J. Hered.* 100 (1), 11–24. <https://doi.org/10.1093/jhered/esn039>.
- Ugland, K.I., Stromnes, E., Berland, B., Aspholm, P.E., 2004. Growth, fecundity and sex ratio of adult whaleworm (*Anisakis simplex*; Nematoda, Ascaridoidea, Anisakidae) in three whale species from the North-East Atlantic. *Parasitol. Res.* 92: 484-489.
- Umehara, A., Kawakami, Y., Matsui, T., Araki, J., Uchida, A., 2006. Molecular identification of *Anisakis simplex* sensu stricto and *Anisakis pegreffii* (Nematoda: Anisakidae) from fish and

917 cetaceans in Japanese waters. Parasitol. Int. 55, 267-271.

1
2
3 <https://doi.org/10.1016/j.parint.2006.07.001>.

4
5
6
7
8
9
10 Valentini, A., Mattiucci, S., Bondanelli, P., Webb, S.C., Mignucci-Giannone, A., Colom, Llavina,
11 M.M., Nascetti, G., 2006. Genetic relationships among *Anisakis* species (Nematoda:
12 Anisakidae) inferred from mitochondrial *cox2* sequences, and comparison with allozyme data.
13 J. Parasitol. 92, 156-166. <https://doi.org/10.1645/GE-3504.1>.

14
15
16
17
18
19
20
21 Van Oosterhout, C., Hutchinson, W.F., Wills, D.P., Shipley, P., 2004. MICRO- CHECKER:
22 software for identifying and correcting genotyping errors in microsatellite data. Mol. Ecol.
23 Notes 4, 535-538. <https://doi.org/10.1111/j.1471-8286.2004.00684.x>.

24
25
26
27
28 Weir, B.C., Cockerham, C.C., 1984. Estimating F-statistics for the analysis of population structure.
29 Evolution 38, 1358–1370. <https://doi.org/10.2307/2408641>.

Table 1
Host species, life-history stage and sex of *Anisakis* specimens (*N*) collected in definitive and intermediate/paratenic hosts from off the New Zealand coast, genotyped at nuclear (*nas10* nDNA, SSRs and mitochondrial gene (mtDNA *cox2*) loci.

Hosts		<i>N</i>	<i>N</i> _A ♂♂	<i>N</i> _A ♀♀	<i>N</i> _{L4}	<i>N</i> _{L3}	<i>N</i> <i>nas10</i> nDNA	<i>N</i> SSRs	<i>N</i> mtDNA <i>cox2</i>
Definitive Hosts									
Delphinidae	<i>Cephalorhynchus hectori</i>	3	1	2	-	-	3	3	3
	<i>Delphinus delphis</i>	28	9	13	6	-	28	28	28
	<i>Globicephala melas</i>	191	52	79	60	-	191	191	191
	<i>Grampus griseus</i>	3	-	-	3	-	3	3	3
Intermediate/Paratenic Hosts									
Oreosomatidae	<i>Allocyttus</i> sp.	9	-	-	-	9	9	9	9
Centrolophidae	<i>Hyperoglyphe antarctica</i>	9	-	-	-	9	9	9	9
Trachichthyidae	<i>Hoplostethus atlanticus</i>	24	-	-	-	24	24	24	24
Merlucciidae	<i>Macruronus novaezealandiae</i>	25	-	-	-	25	25	25	25
Moridae	<i>Mora moro</i>	19	-	-	-	19	19	19	19
	<i>Pseudophycis bachus</i>	12	-	-	-	12	12	12	12
Gempylidae	<i>Rexea solandri</i>	10	-	-	-	10	10	10	10
Carangidae	<i>Trachurus declivis</i>	11	-	-	-	11	11	11	11
Tot:		344	62	94	69	119	344	344	344

Table 2
Assignment of the *Anisakis* specimens tested in the present study to the parental taxa *Anisakis pegreffii* (Ap) or *A. berlandi* (Ab), or to mixed ancestry between them (Heterozygote), according to their genotypes observed at nuclear (i.e. *nas10* gene and SSRs loci) and mitochondrial (mtDNA *cox2*) markers.

Host species	N	<i>nas10</i> nDNA			SSRs			mtDNA <i>cox2</i>	
		Ap	Ab	Heterozygote	Ap	Ab	Heterozygote	Ap	Ab
Definitive Hosts	$N_{A, L4}$								
<i>Cephalorhynchus hectori</i>	3	3	-	-	3	-	-	3	-
<i>Delphinus delphis</i>	28	28	-	-	28	-	-	28	-
<i>Globicephala melas</i>	191	137	53	1	137	53	1	137	54
<i>Grampus griseus</i>	3	3	-	-	3	-	-	3	-
Intermediate/Paratenic Hosts	N_{L3}								
<i>Allocyttus</i> sp.	9	-	9	-	-	9	-	-	9
<i>Hyperoglyphe antarctica</i>	9	-	9	-	-	9	-	-	9
<i>Hoplostethus atlanticus</i>	24	-	24	-	-	24	-	-	24
<i>Macruronus novaezelandiae</i>	25	3	22	-	3	22	-	3	22
<i>Mora moro</i>	19	8	11	-	8	11	-	8	11
<i>Pseudophycis bachus</i>	12	9	3	-	9	3	-	9	3
<i>Rexea solandri</i>	10	8	2	-	8	2	-	8	2
<i>Trachurus declivis</i>	11	9	-	2	9	-	2	9	2
Tot:	344	208	133	3	208	133	3	208	136

Table 3

Genetic diversity at nine microsatellite loci of adult and larval specimens in populations of *A. pegreffii* and *A. berlandi* analysed in the present study. *Anisl 2* and *Anisl 15* were excluded from the estimates because they are monomorphic in both the species. Similarly, no estimates were possible for the loci *Anisl 00314* and *Anisl 22* loci because they are monomorphic in both larvae and adults of *A. berlandi*, as well as in *Anisl 10535* locus because it was monomorphic in adult and L4 specimens.

N, number of genotyped specimens at each locus; *Ho*, observed heterozygosity; *He*, expected heterozygosity; *A*, number of alleles detected at each locus; *p*, significance ($p < 0.05$) of the deviation from HWE expectation. *** $p < 0.001$.

		<i>A. pegreffii</i>		<i>A. berlandi</i>	
		A+L4	L3	A+L4	L3
Locus					
<i>Anisl 00185</i>	<i>N</i>	171	37	53	80
	<i>Ho</i>	0.75	0.76	0.74	0.77
	<i>He</i>	0.81	0.79	0.83	0.84
	<i>p</i>	0.61	0.73	0.51	0.33
	<i>A</i>	9	7	11	9
<i>Anisl 00314</i>	<i>N</i>	171	37	53	80
	<i>Ho</i>	0.45	0.35	-	-
	<i>He</i>	0.75	0.74	-	-
	<i>p</i>	***	***	-	-
	<i>A</i>	8	6	1	1
<i>Anisl 10535</i>	<i>N</i>	171	37	53	80
	<i>Ho</i>	0.73	0.65	-	0.01
	<i>He</i>	0.76	0.79	-	0.01
	<i>p</i>	0.45	0.55	-	1.00
	<i>A</i>	9	8	1	2
<i>Anisl 05784</i>	<i>N</i>	171	37	53	80
	<i>Ho</i>	0.81	0.81	0.07	0.02
	<i>He</i>	0.79	0.77	0.07	0.02
	<i>p</i>	0.09	0.31	1.00	1.00
	<i>A</i>	12	9	2	2
<i>Anisl 08059</i>	<i>N</i>	171	37	53	80
	<i>Ho</i>	0.83	0.86	0.41	0.30
	<i>He</i>	0.86	0.87	0.36	0.26
	<i>p</i>	0.05	0.66	0.84	0.45
	<i>A</i>	14	12	4	3
<i>Anisl 00875</i>	<i>N</i>	171	37	53	80
	<i>Ho</i>	0.44	0.49	0.07	0.11
	<i>He</i>	0.49	0.42	0.07	0.11
	<i>p</i>	0.28	1.00	1.00	1.00
	<i>A</i>	8	6	3	4
<i>Anisl 7</i>	<i>N</i>	171	37	53	80
	<i>Ho</i>	0.21	0.13	0.56	0.47
	<i>He</i>	0.25	0.33	0.91	0.92
	<i>p</i>	***	***	***	***
	<i>A</i>	4	4	21	23
<i>Anisl 4</i>	<i>N</i>	171	37	53	80
	<i>Ho</i>	0.53	0.43	0.43	0.35
	<i>He</i>	0.82	0.84	0.82	0.85
	<i>p</i>	***	***	***	***
	<i>A</i>	12	12	14	15
<i>Anisl 22</i>	<i>N</i>	171	37	53	80
	<i>Ho</i>	0.41	0.32	-	-
	<i>He</i>	0.61	0.62	-	-
	<i>p</i>	***	***	-	-
	<i>A</i>	8	5	1	1

Table 4

Genetic diversity at the four sex-linked loci (i.e. *Anisl 00314*, *Anisl 7*, *Anisl 4* and *Anisl 22*) in *A. pegreffii* and *A. berlandi*, estimated in adult female specimens. Estimates at the *Anisl 00314* and *Anisl 22* loci were not possible because they are monomorphic in *A. berlandi*. *N*= number of genotyped specimens at each locus; *Ho*, observed heterozygosity; *He*, expected heterozygosity; *p*, significance ($p < 0.05$) of the deviation from HWE expectation. *** $p < 0.001$.

		<i>A. pegreffii</i> ♀♀	<i>A. berlandi</i> ♀♀
Locus			
<i>Anisl 00314</i>	<i>N</i>	71	22
	<i>Ho</i>	0.67	-
	<i>He</i>	0.77	-
	<i>p</i>	0.17	-
<i>Anisl 7</i>	<i>N</i>	71	22
	<i>Ho</i>	0.30	1.00
	<i>He</i>	0.30	0.87
	<i>p</i>	0.67	0.51
<i>Anisl 4</i>	<i>N</i>	71	22
	<i>Ho</i>	0.79	0.77
	<i>He</i>	0.85	0.87
	<i>p</i>	0.63	0.60
<i>Anisl 22</i>	<i>N</i>	71	22
	<i>Ho</i>	0.61	-
	<i>He</i>	0.63	-
	<i>p</i>	0.27	-

Table 5

Allele frequencies observed at eleven microsatellite loci of *A. pegreffii* and *A. berlandi* here studied. As regards sex-linked loci (*; i.e. *Anisl 00314*, *Anisl 7*, *Anisl 4*, *Anisl 22*), the most reliable estimate of allele frequencies was calculated according to the sex-linked genetic model estimate, assuming: (i) the hemizyosity of the males at that locus in the two *Anisakis* species; (ii) their adult female counterparts, as biallelic at the sex-linked loci.

Locus	Alleles	<i>A. pegreffii</i>		<i>A. berlandi</i>	
		A + L4	L3	A + L4	L3
<i>Anisl 00185</i>	182	0.01	-	0.01	-
	185	-	-	0.01	-
	188	0.02	0.05	0.03	0.01
	191	0.06	0.04	0.07	0.15
	194	0.21	0.27	0.16	0.17
	197	0.21	0.26	0.21	0.19
	200	0.17	0.26	0.29	0.21
	203	0.25	0.11	0.10	0.12
	206	0.05	0.01	0.07	0.12
	209	0.02	-	0.03	0.01
	212	-	-	0.02	0.02
<i>Anisl 00314*</i>	96	0.05	-	-	-
	100	0.28	-	1.00	-
	104	0.32	-	-	-
	108	0.25	-	-	-
	112	0.06	-	-	-
	116	0.01	-	-	-
	120	0.02	-	-	-
	124	0.01	-	-	-
<i>Anisl 10535</i>	125	0.01	-	-	-
	128	0.01	0.01	1.00	0.99
	131	0.01	0.01	-	0.01
	134	0.19	0.20	-	-
	137	0.21	0.18	-	-
	140	0.27	0.27	-	-
	143	0.28	0.28	-	-
	146	0.01	0.04	-	-
	149	0.01	-	-	-
	152	-	0.01	-	-
<i>Anisl 05784</i>	66	-	0.01	-	-
	69	0.01	-	0.96	0.99
	72	0.01	-	0.04	0.01
	75	0.01	0.05	-	-
	78	0.05	0.02	-	-
	81	0.03	0.04	-	-
	84	0.07	0.09	-	-
	87	0.25	0.23	-	-
	90	0.32	0.40	-	-
	93	0.18	0.12	-	-
	96	0.05	0.04	-	-
	99	0.01	-	-	-
	102	0.01	-	-	-

(continues)

Table 5
continued

Locus	Alleles	<i>A. pegreffii</i>		<i>A. berlandi</i>	
		A + L4	L3	A + L4	L3
<i>Anisl 08059</i>	82	0.02	0.01	0.01	-
	86	0.19	0.19	0.77	0.85
	90	0.05	0.06	0.20	0.14
	94	0.12	0.19	0.02	0.01
	98	0.17	0.16	-	-
	102	0.20	0.16	-	-
	106	0.11	0.08	-	-
	110	0.04	0.05	-	-
	114	0.05	0.02	-	-
	118	0.01	0.01	-	-
	122	0.01	0.05	-	-
	126	0.01	0.02	-	-
	130	0.01	-	-	-
	134	0.01	-	-	-
<i>Anisl 00875</i>	142	-	-	-	0.01
	145	0.02	-	-	0.01
	148	0.02	-	0.96	0.94
	151	0.04	0.08	0.03	0.04
	154	0.03	0.04	0.01	-
	157	0.68	0.76	-	-
	160	0.15	0.08	-	-
	163	0.05	0.03	-	-
	166	0.01	-	-	-
	169	-	0.01	-	-
<i>Anisl 7*</i>	216	0.02	-	-	-
	219	0.07	-	-	-
	222	0.84	-	-	-
	225	0.07	-	0.01	-
	228	-	-	0.01	-
	237	-	-	0.01	-
	243	-	-	0.01	-
	252	-	-	0.01	-
	255	-	-	0.01	-
	258	-	-	0.06	-
	261	-	-	0.07	-
	264	-	-	0.05	-
	267	-	-	0.19	-
	270	-	-	0.19	-
	273	-	-	0.14	-
	276	-	-	0.04	-
	279	-	-	0.04	-
	282	-	-	0.08	-
	285	-	-	0.01	-
	288	-	-	0.02	-
	291	-	-	0.01	-
	294	-	-	0.02	-
	297	-	-	0.01	-
	306	-	-	0.01	-

(continues)

Table 5.
Continued

Locus	Alleles	<i>A. pegreffii</i>		<i>A. berlandi</i>	
		A + L4	L3	A + L4	L3
<i>Anisl 4*</i>	127	0.04	-	0.13	-
	130	0.06	-	0.02	-
	133	0.31	-	0.02	-
	136	0.12	-	-	-
	139	0.16	-	0.37	-
	142	0.16	-	0.08	-
	145	0.07	-	0.04	-
	148	0.02	-	-	-
	151	0.02	-	0.02	-
	154	0.02	-	0.04	-
	157	0.01	-	0.05	-
	160	-	-	0.13	-
	163	-	-	0.04	-
	166	-	-	0.02	-
	169	-	-	0.02	-
	172	-	-	0.02	-
	175	0.01	-	-	-
<i>Anisl 22*</i>	179	0.02	-	1.00	-
	182	0.04	-	-	-
	185	0.45	-	-	-
	188	0.40	-	-	-
	191	0.06	-	-	-
	194	0.01	-	-	-
	197	0.01	-	-	-
	200	0.01	-	-	-
<i>Anisl 15</i>	231	1.00	1.00	-	-
	235	-	-	1.00	1.00
<i>Anisl 2</i>	245	-	-	1.00	1.00
	248	1.00	1.00	-	-

Table 6.

Analysis of molecular variance (AMOVA) between and within adult and L4 stages group vs L3 stages group, of the two species *A. pegreffii* (a) and *A. berlandi* (b), using genetic data sets from seven SSRs loci (*Anisl 00185*, *Anisl 10535*, *Anisl 05784*, *Anisl 08059*, *Anisl 00875*, *Anisl 15* and *Anisl 2*). The *p* values to the fixation indices are reported as ****p*<0.001, * *p*<0.05. n.s., not significant.

a)

<i>Anisakis pegreffii</i>					
Source of variation	d.f.	Sum of squares	Variance components	% of variation	Fixation indices
Among populations	1	2.413	0.004	0.22	<i>Fst</i> = 0.002 n.s.
Among individuals within populations	185	356.785	0.075	4.05	<i>FIS</i> = 0.040*
Within individuals	187	332.500	1.778	95.73	<i>FIT</i> = 0.043*
Total	373	691.698	1.857		

b)

<i>Anisakis berlandi</i>					
Source of variation	d.f.	Sum of squares	Variance components	% of variation	Fixation indices
Among populations	1	0.889	0.002	0.28	<i>Fst</i> = 0.003 n.s.
Among individuals within populations	131	86.341	0.016	2.42	<i>FIS</i> = 0.024 n.s.
Within individuals	133	83.500	0.628	97.30	<i>FIT</i> = 0.027 n.s.
Total	265	170.729	0.645		

	240		250		260																									
<i>A. pegreffii</i>		T	A	T	G	A	T	T	G	A	T	G	C	G	T	T	A	A	T	T	C	T	T	T	C	T	A	C		
<i>A. berlandi</i>		.	.	.	.	.	A	.	.	.	.	.	.	C	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	
Heterozygote		.	.	.	.	.	.	.	.	.	.	.	S	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.		





Figure 2

[Click here to access/download;Figure\(s\);Fig2.tif](#)








Figure 4

[Click here to access/download;Figure\(s\);Fig4.tif](#)





Click here to access/download
Supplementary Material
Suppl.Fig1.tif



Conflict of Interest and Authorship Conformation Form

Please check the following as appropriate:

- ☐ All authors have participated in (a) conception and design, or analysis and interpretation of the data; (b) drafting the article or revising it critically for important intellectual content; and (c) approval of the final version.
- ☐ This manuscript has not been submitted to, nor is under review at, another journal or other publishing venue.
- ☐ The authors have no affiliation with any organization with a direct or indirect financial interest in the subject matter discussed in the manuscript
- ☐ The following authors have affiliations with organizations with direct or indirect financial interest in the subject matter discussed in the manuscript:

Author's name	Affiliation
Eleonora Bello	Tuscia University
Marialetizia Palomba	Sapienza University
Stephen C. Webb	Cawthron Institute
Michela Paoletti	Tuscia University
Paolo Cipriani	Institute of Marine Research
Giuseppe Nascetti	Tuscia University
Simonetta Mattiucci	Sapienza University